



Influence of Different Pesticides on Chemical, Biochemical and Yield Parameters of Brinjal (*Solanum Melongena L*)

S. B. Gaikwad¹, M. B. Chetti² and D. I. Jirali³

Email.id. swati5688@yahoo.com

Dr. Gaikwad Swati Bhimraoji (Ph.D crop Physiology) UAS, Dharwad, Karnataka, India, 560005 Postal address: C/o, Usha Thakare, Isaji Lay Out, Ward number 6, Near Kesarimal Kanya School, Wardha 442001, Maharashtra, India, Mobile number: 8867636330

Dr. M. B. Chetti (Ph.D, Plant physiology) ADG-ICAR, pusa New Delhi, Human Resources Department, India, email.id mbchetti_uas@rediffmail.com

Dr. D. I. Jirali, Professor of crop Physiology, Dept of CPH, UAS Dharwad, Karnataka, email.id. dijirali_uasd@rediffmail.com

Abstract:

Pesticides have become an essential part of agricultural and horticultural practices. There is continuous development of new pesticides to address specific requirements (insect/pest control). New pesticides with specific mode of action, combination to two or more pesticides for enhanced bio-efficacy, new innovation in the field of slow and sustained delivery of chemical are making the pesticides influence positively in growth and development of plant. In present investigation, all four pesticides used at recommended dosage suggested by agricultural experts. These pesticides are most preferred by farmers to control fruit and shoot borer on brinjal. A study on the influence of pesticides persistence on biochemical and yield parameters of brinjal was conducted during *kharif* and *rabi* 2013-14 and 14-15 with the objective of physiological responses of different varieties of brinjal (Malapur local-V₁, Kalpataru-V₂, Manjula-V₃, Manjari-V₄) sprayed with four different pesticides (thiodicarb-P₁, spinosad-P₂, profenophos-P₃ and chlorantraniliprole-P₄). The P₀ treatment considered as a non-sprayed control. The experiment was laid out in factorial randomized block design with three replication. The pesticides sprayed at recommended dose (thiodicarb 75WP@1g⁻¹, spinosad 45 SC @0.1 ml⁻¹ profenophos 50EC@2ml⁻¹ and chlorantraniliprole 20 SC @ 0.5 ml⁻¹) suggested by agricultural experts. Results indicated highly significant differences between the varieties and pesticides treatments at all the stages and in both the seasons. The interaction between the varieties and pesticide treatments was also significant at all the stages. The values of chlorophyll content successive increased from 70 to 90 DAT and decreases from 115 and 125 DAT in both the seasons. The magnesium content in fruits was recorded highest at 90 DAT followed by 80 DAT irrespective of varieties and pesticides. It is clear from the data that (V₄P₃) indicated maximum magnesium content recorded followed by V₄P₄ compared to other interaction. Total marketable yield noticed maximum in Manjari followed by Manjula and Kalpataru while least was in Malapur local. The profenophos treatment observed with highest marketable yield over the chlorantraniliprole. Spinosad and thiodicarb treatment. Among all the interaction V₄P₃ (6.12 kg plant⁻¹) recorded maximum marketable yield and minimum was in V₁P₀ interaction.

Key words: Pesticides, residues, chlorophyll, magnesium and brinjal

Introduction

Pesticides are one of the most essential inputs in modern agriculture and horticulture for insuring food security. Among various strategies adapted to combat pest of brinjal, pesticides from the first choice of farmers. Most of the pesticides that are used against pest on brinjal belong to organophosphates. But the indiscriminate use of pesticides, has led to various secondary effects on plant growth and development. Some pesticides remain persistent and take a long time to move into the environment. The brinjal, *solanum melongena L.* are an important vegetable crop grown throughout the year in India. However, brinjal heavily suffers from shoot and fruit borer from seedling stage to fruiting, which reduce not only the yield but also quality of fruits. The pesticides stress at high concentration of pesticides adversely affect the chlorophyll stability, membrane stability index and relative water content (Nasrabadi and Dhumal, 2014). Pesticides play an important role in plant protection, but their excess and continuous use has harmful effect on growth, development, yield

and grown in crop plant. In present investigation the organophosphate group of pesticides (profenophos) play an important role in controlling shoot and fruit borer followed by chlorantraniliprole at recommended dosage. At recommended dosage of pesticides with minimum frequency results increase in chlorophyll content of leaves and magnesium content of fruits.

Materials and methods

Chlorophyll content in the leaves of different genotypes of brinjal leaves was determined by using dimethyl sulphoxide (DMSO) as given by Shoaf and Loum (1976). All pesticide residues analysed by liquid chromatography and tandem mass spectroscopy and calculated in mg kg⁻¹ of fruit weight at different days after spraying. Magnesium in digested samples was determined by EDTA titration. The marketable fruit yield per plant was calculated by substituting the fruit weights of borer infected fruits and unmarketable fruits of each picking and expressed in gm per plant

Results

Total chlorophyll content ($\mu\text{g g}^{-1}$)

The results of total chlorophyll content of brinjal leaf due to application of pesticides given in Table 1 (a) indicated significant differences between the varieties and pesticide treatments at all stages and in both the seasons. The interaction between varieties and pesticide treatments was also significant at all the stages. The pooled data showed that V₄ noted maximum value (659.4 $\mu\text{g g}^{-1}$) followed by V₂ and V₃ and least was in V₁ at all stages. While, among the pesticide treatments, it was maximum in P₃ followed by P₄, P₂, P₁ and the minimum was in P₀. Among different stages, it was maximum at 90 DAT followed by 115 and 80 DAT; while, least was at 125 DAT. It increased from 70 to 90 DAT in all treatments and declined slowly at 125 DAT. A maximum chlorophyll content (970 $\mu\text{g g}^{-1}$) was recorded in V₄P₃ while, the lowest content was recorded in V₁P₀ followed by V₁P₁ and V₁P₂ at all the stages. The treatment V₁P₄ did not differ significantly with V₂P₀ at 70, 80, 90 and at 115. Similar results were obtained in *rabi* as that of *kharif*. Data presented in Table 1 (b) showed maximum total chlorophyll content in V₄P₃ (961.7 $\mu\text{g g}^{-1}$) among all interactions. The treatment V₄P₁ did not differ significantly with V₄P₀ at 125 DAT.

Magnesium content in fruits (mg 100 g⁻¹)

The data on magnesium content presented in Table 2 (a) indicated significant differences between the varieties and pesticide treatments at all stages and in both the seasons. The pooled data showed that V₄ recorded maximum value (11.70 mg 100 g⁻¹ at 90 DAT) followed by V₂ and V₃ and least was in V₁. While, among the pesticide treatments, it was maximum in P₃ followed by P₄ and minimum in P₀. Among different stages, it was maximum at 90 DAT followed by 80 and 115 DAT; while least was at 125 DAT. A maximum magnesium content was noted in P₃ (9.90 mg 100 g⁻¹).

Interaction between varieties and pesticides revealed significant differences at 80, 90 and 125 DAT. While there were no significant differences at 70 and 115 DAT. It is clear from the data that V₄P₃ indicated maximum value at all the stages compared to other treatment combinations. The maximum magnesium content was recorded at 90 DAT followed by 80 DAT irrespective of varieties and pesticides. However, significantly lower magnesium content was recorded in V₁P₀ at all the stages. The magnesium content increased from 70 to 90 DAT in all treatments and declined slowly thereafter until 125 DAT. A maximum magnesium content of 17.65 mg 100 g⁻¹ was recorded in V₄P₃ at 90 DAT;

while the lowest magnesium content (10.32 mg 100 g⁻¹) was recorded in V₁P₀ at 70 DAT. The data on magnesium content presented in Table 29 (b) indicated significant differences between the varieties and pesticide treatments at all stages and in both the *rabi* seasons. The interaction between varieties and pesticide treatments was also significant at all the stages in both the seasons. Similar results were obtained in *rabi* as that of *kharif* with respect to in varieties and pesticide treatments. Maximum magnesium content was observed in V₄P₃ (17.65 mg 100 g⁻¹) among all interactions followed by V₄P₃ and V₂P₃ at all the stages. The minimum magnesium content was noticed in V₁P₀ (10.31 mg 100 g⁻¹) at 70 DAT.

Marketable fruit yield

The highest marketable fruit yield was recorded in V₄ at 90 DAT among all the varieties. V₁ produce significantly lowest yield compared to other varieties at all the stages. It was calculated highest from P₃ which was closely followed by P₄, P₂ and P₁ whereas lowest in P₀ (non-sprayed control). Pesticide treatments, varieties and their interactions differed significantly among themselves at all the stages.

It is clear from the data in Table 3(a) that V₄P₃ recorded maximum value at all the stages compared to other treatment combinations. The highest marketable fruit yield was observed at 90 DAT followed by 80 DAT; irrespective of varieties and pesticides. However, significantly lowest was calculated from V₁P₀ at all the stages. It is clear from the data that highest total marketable yield (7.09 kg) was recorded in V₄P₃ while the lowest in V₁P₀ (1.88 kg plant⁻¹). The treatment V₂P₁ did not differ significantly with V₂P₂ at 90 DAT. Similarly, the treatments V₁P₁ and V₃P₃; at 90 and V₂P₂ and V₂P₃; V₄P₃ and V₄P₄; V₁P₃ and V₁P₄ at 125 DAT did not differ significantly among themselves. Pooled data recorded showed significantly differ marketable fruit yield in *rabi* at all the stages in varieties, pesticide treatments and their interactions respectively. Results obtained was followed same trend as that of *kharif*. The highest marketable yield was noticed in V₄P₃ (1.43 kg plant⁻¹) at 90 DAT; while lowest in V₁P₀ (0.21 kg) at 80 DAT.

Discussion

Data presented in Tables 1(a) and 1(b) revealed a significant differences between the varieties, pesticide treatments and their interactions at different days after transplanting and in both the seasons (*kharif* 2013-14 and 14-15 and *rabi* 2013-14 and 14-15) after fifth sprays in chlorophyll content. There was a increase in chlorophyll content during the growth period 70-

90 DAT appeared in treated brinjal leaves. This results add further support to the changes in photosynthetic activity and/or alteration in chloroplast. In addition, the pesticides act as a chemical stressor which may interrupt the electron transport activity in PSII, cytochrome b6/f or PSI or may alter the structure of the chloroplast, or inhibit the calvin cycle (Lichtenthaler 1987). Generally, there might be different responses according to the concentrations and frequency of spraying of the pesticide. It also depends on type of pesticides and its period of persistence.

Previously, an increase in the chlorophyll content was observed in *S. tuberosum* by deltamethrin (Fidalgo *et al.*, 2000) and in *H. esculentus* and *C. annuum* by Topsin (Ahmed and Siddiqui, 1995); whereas reduction by butachlor on rice (Bhattacharya *et al.*, 2001) was recorded. He also observed carbofuran, butachlor and carbendazim on chlorophyll and ultra structural alternation to *Solanum tuberosum* chloroplasts in *Solanum melongena*. Effects of different doses of endosulfan on total chlorophyll contents of cabbage leaf on 1, 7, 14, 21, 28 and 35 days after application (DAA) increases of 25.64, 23.81 and 27.50 per cent respectively in chlorophyll content were found on 35th DAA with respect to recommended dose of 0.844 L ha⁻¹ (Ashrafi and Pandit, 2015). In this work, chlorophyll content is highly relevant to well determine the effect of thiodicarb, spinosad and profenophos and chlorantraniliprole on photosynthesis of four different varieties of brinjal. For this, all measurements were preferred on a young leaves. This study confirms that at recommended dose of profenophos, rynaxypyr and spinosad and thiodicarb respectively increases the total chlorophyll content in brinjal leaves (Manjari, Kalpataru, Manjula and Malapur local). The variety Manjari noticed with maximum chlorophyll content at 90 DAT with respect to profenophos treatments (V₄P₃) followed by V₂P₃ and V₄P₄ might be due to persistence of pesticide of profenophos and chlorantraniliprole, due to low dose of pesticides. The lowest chlorophyll content observation in thiodicarb followed by spinosad pesticide treatments probably due to its easy volatilization or degradation rate.

In contrast to this study the researchers like Abbas *et al.* (1994) claimed that phenolics inhibit CO₂ dependant O₂ evolution in intact chloroplast. Singh *et al.* (1994) reported that phenolic compounds inhibit photosynthesis in intact plants, which results in reduced growth and yield. The decrease in chlorophylls and photosynthetic rate may affect the plant

growth. The pesticides may affect some biochemical composition of plant. The organophosphorus compounds used in agriculture usually alter the chemical composition and nutritive value of plant product. Despite the simplicity in structure and simple mode of action, the cyanophos (phosphorothioate) is typical xenobiotics (Neumann and Peter, 1987) and has an adverse effect on the plant. The common mechanism of its toxic action is inhibition of biological pathways such as photosynthesis and mitochondrial electron transport. Plant productivity depends on the conversion of light energy into stable chemical energy. If the photosynthetic apparatus is inhibited by environmental contaminants, changes in plant cell physiology, growth and biomass yield are inevitable. As well it has been shown that inhibition of photosynthesis is a reliable assay of the potential toxicity and xenobiotics contaminants towards plants (Mallakin *et al.*, 2002). Therefore, in the present study, recommended dose fixed by agricultural experts with the logic of spraying less often and using fewer and better selected pesticides only when needed can reduce the pesticides stress in brinjal plants by minimizing residues.

At low dose of imidacloprid significantly promoted the uptake of P and K by rice plants (Quiet *et al.*, 2004). The presence of pesticides residues like chlorosulfuron, dclorop, haloxyfopfluazifop have shown some specific effects on the micronutrient transport system and the plasma membrane of the root cells (Shimabukuro, 1990; Shimabukuro & Hoffer, 1994; Wright, 1994) which in turn affect the cations uptake including Zn, Cu and Mn (Tester, 1990). Photosynthesis is a more general measurement of carbon dioxide intake and fixation (sugar production). Both processes are interrelated and are directly related to plant growth and productivity. The higher value of transpiration occurred on pesticides-free plants. Photosynthesis was also higher in the untreated check, suggesting that all pesticide treatments at higher rate adversely influenced both stomatal opening and overall photosynthetic rates and that the effect remained for at least one week. Reductions of lettuce head weight and diameter in plants treated weekly with methyl-parathion indicate the cumulative effect on yield that results from pesticide inhibition of physiological processes. Since no differences were observed between the photosynthesis and transpiration rates of plants treated in daylight or darkness, it may be assumed that pesticides uptake by the plant is independent of stomatal opening.

The pesticides used in experiments i.e. thiodicarb, spinosad, profenophos and chlorantraniliprole at recommended doses are from different classes, differ in their effects on plants. Additionally, the rates, number and timing of applications may alter a compound effect upon the plants, for either or short time or several days. Efficiency of pesticides treatment to reduce per cent fruit borer infestations may increase the yield from 70 to 90 DAT (10 to 20 days after fifth spraying) while, slightly decline in yield during both *kharif* and *rabi* due to increase in pest infestations after 90 DAT. Results obtained on the influence of different pesticides in different growing seasons on yield of brinjal varieties with certain environmental parameter, when relative effectiveness of pesticides in reducing pest populations is considered, the later compound are most desirable. As more data are obtained on the effect of pesticides and their compounds on photosynthesis and consequently on yield may become essential. Results indicate that the dual effect of profenophos treatments on different varieties, by measuring changes in photosynthesis (alteration in chlorophyll content) and most efficient pesticides among all pesticide treatments reduced per cent fruit borer infestations from brinjal plant.

Conclusion

The total chlorophyll content in brinjal leaves can be very useful in detecting the effects of different pesticides on the plant physiological processes caused by recommended dose of all four pesticides and its gave us an alternative view on the action mechanism of all four pesticides on photosynthesis of treated brinjal. Present study showed that there is a difference between all four varieties treated with different pesticides, and variety Manjari was more tolerant followed by Kalpataru, Manjula and most sensitive variety Malapur local to pesticide exposure. Results find that the difference between all four pesticides treatments in responses to varieties probably due to accumulation of ROS increasing rate vary with varieties which contribute to oxidative stress that touches photosynthesis and the biosynthesis of photosynthetic pigments. The profenophos followed by chlorantraniliprole, spinosad and thiodicarb significantly promoted the uptake of phosphorus, sulphur, magnesium and calcium by brinjal plant. Thus, it is difficult to make any generalization about the impact of pesticides on brinjal plants. The present study demonstrates that there were no inhibitory effects on the growth, photosynthetic activity and yield of all brinjal varieties at low dosage and interestingly profenophos followed by chlorantraniliprole,

spinosad and thiodicarb support the plant activity over the non-sprayed controls. It can be concluded that recommended dose of profenophos after 20 DAS (90 DAT) is beneficial for the growth, photosynthetic pigments and yield of brinjal varieties specifically in Manjari (V4) among all varieties examined.

References

- Abbas, H. K., Duke, S. O., Paul, R. N., Riley, R. T. and Tanaka, T., 1994, AAL-toxin, a potent natural herbicide with disrupting of lipid metabolism of plants. *Pesticide Sci*, 43: 181-187.
- Ahmed, S. and Siddiqui, Z. S., 1995, Effect of topsin fungicide (methylthiophenate) on chlorophyll, protein and phenol contents of *Hibiscus esculentus* and *Capsicum annum*. *Pakistan J. Botany*, 27: 175-178.
- Ashrafi, A. A. and Pandit, G. K., 2015, Variation in the formation of some biomolecules in cabbage (*Brassica oleracea* var. capitata) leaf induced by endosulphon. *The Bioscan*, 10 (2) : 539-543.
- Bhattacharya, A., Pandit, G. K., Chakraborti, P. and Roy, D., 2001, Effect of carbofuran, butachlor and carbendazim on chlorophyll and carbohydrate contents of two summer rice (*Oryza sativa* L.) cultivars at panicle emergence. *Crop Res.*, 21: 15-19.
- Fidalgo, F., Santos, I. and Salema, R., 2000, Nutritional value of potato tubers from field grown plants treated with deltamethrin. *Potato Res.*, 43 : 43.
- Lichtenthaler, H. K., 1987, Chlorophylls and carotenoid pigments of photosynthetic biomembranes. *Methods of Enzymology*, 148: 350-366.
- Mallakin, A., Babu, T. S., Dixon, D. G. and Greenberg, B. M., 2002, Sites of toxicity of specific photooxidation products of anthracene to higher plants: Inhibition of photosynthetic activity and electron transport in *Lemna gibba* L. G-3 (Duckweed). *Environ. Toxicol.*, 17 (5) : 462-471. 12.
- Nasrabadi, M., Ghayal, N. and Dhumal, K. N. 2011. Effect of chloropyrifos and malathione on antioxidant enzymes in tomato and brinjal. *Int. J. Pharma Biosci.*, 2 (2) : 202-208.
- Neumann, I. and Peter, H. H., 1987, Insecticidal organophosphates: nature made them first. *Experimentia*, 43: 1235-1237.
- Qui, H. M., Wu, J. C., Yang, G. Q., Dong, B. and Li., D. H., 2004, Changes in uptaking function of rice root to nitrogen, phosphorus and potassium under brown planthopper, *Nilparvata lugens* (Stål) (Homoptera:

Delphacidae) and pesticides stresses. *Crop Prot.*, 10; 1016.

Shimabukuro, R. H. and Hoffer, B. L., 1994, Induction of ethylene as an indicator of senescence in the mode of action of diclofop-methyl. *Pestic. Biochem. Physiol.*, 54 (2) : 146-158.

Shimabukuro, R. H., 1990, Biosciences research laboratory, USDA, Agricultural Research services, state university stolon, Fargo, North Dakota 58105-5674, USA.

Shoaf, T. W. and Loum, B. W., 1976, Improved extraction of *chlorophyll a* and *b* from algae using Dimethyl sulfoxide. *Oceanography*, 21: 926-928.

Tester, M., 1990, Plant Ion Channels: whole-cell and single channels studies. *NewPhytol.*, 114: 305-340

Wright, J. P., 1994, Use of membrane potential measurements to study mode of action of diclofop-methyl. *Weed Sci.*, 42 (2) : 285-292.

Table 1(a). Influence of pesticides on total chlorophyll content ($\mu\text{g g}^{-1}$) of brinjal in different growing seasons

Treatments	Pooled data (kharif 2013-14 & 14-15)				
	70	80	90	115	125
Varieties (V)					
V ₁	441.1	464.5	509.0	492.2	421.9
V ₂	563.2	586.1	632.6	611.5	547.1
V ₃	563.1	584.4	629.1	614.5	544.8
V ₄	599.5	621.0	664.6	643.2	572.0
Mean	541.7	564.0	608.8	590.4	521.5
S.E.m±	0.29	0.29	0.29	0.28	0.29
LSD @ 5%	0.83	0.83	0.83	0.81	0.83
Pesticides (P)					
P ₀	391.2	409.4	445.4	428.6	374.8
P ₁	413.2	430.4	466.4	454.6	397.0
P ₂	423.0	440.4	476.1	463.8	405.0
P ₃	481.7	499.4	536.4	517.0	464.6
P ₄	457.9	476.5	511.0	497.5	444.6
Mean	433.4	451.2	487.1	472.3	417.2
S.E.m±	0.30	0.26	0.26	0.25	0.26
LSD @ 5%	0.90	0.74	0.74	0.73	0.74
Interactions (V×P)					
V ₁ P ₀	481.0	511.5	571.5	544.7	441.0
V ₁ P ₁	514.3	544.8	604.8	585.7	494.8
V ₁ P ₂	530.6	561.2	621.2	599.2	501.2
V ₁ P ₃	727.1	757.8	814.7	792.7	704.7
V ₁ P ₄	687.8	721.2	781.2	759.2	671.2
V ₂ P ₀	696.0	726.5	786.5	755.3	669.8
V ₂ P ₁	736.5	767.0	827.0	805.0	712.0
V ₂ P ₂	747.0	777.5	837.5	815.5	727.5
V ₂ P ₃	812.8	843.3	912.8	868.3	794.8
V ₂ P ₄	762.6	793.2	853.2	832.2	743.2
V ₃ P ₀	683.3	713.3	773.3	751.3	663.3
V ₃ P ₁	735.7	764.0	824.0	812.5	710.7
V ₃ P ₂	761.2	787.8	846.2	833.7	734.5
V ₃ P ₃	799.5	826.5	886.5	860.2	772.2
V ₃ P ₄	774.2	804.2	864.2	839.2	751.2
V ₄ P ₀	747.7	777.7	837.7	805.8	724.2
V ₄ P ₁	768.5	793.5	853.5	827.2	729.2
V ₄ P ₂	781.0	809.3	869.3	843.3	736.7
V ₄ P ₃	871.7	901.7	961.7	925.3	825.3
V ₄ P ₄	828.0	858.0	908.2	886.2	798.2
Mean	722.3	752.0	811.7	787.1	695.3
S.E.m±	2.60	2.58	2.58	2.53	2.58
LSD @ 5%	7.44	7.40	7.39	7.25	7.40

Pesticides were detected only at 24 and 96 hr after the spray. No further detection was seen

- V₁ – Malapur local
- V₂ – Kalpataru
- V₃ – Manjula
- V₄ – Manjari
- P₀ – Control (water sprayed)
- P₁ – Thiodicarb 75 WP @ 1g/L
- P₂ – Spinosad 45 SC @ 0.1 ml/L
- P₃ – Profenophos 50 EC @ 2 ml/L
- P₄ – Chlorantraniliprole 20 SC @ 0.5 ml/L

Table 1(b). Influence of pesticides on total chlorophyll content ($\mu\text{g g}^{-1}$) of brinjal in different growing seasons

Pooled data (rabi 2013-14 & 14-15)					
Treatments	DAT				
	70	80	90	115	125
Varieties(V)					
V ₁	440.6	464.4	508.0	483.0	405.5
V ₂	562.6	585.5	630.2	596.8	533.8
V ₃	562.6	585.1	629.8	604.5	538.5
V ₄	591.0	613.0	659.6	634.4	548.4
Mean	539.2	562.0	606.9	579.7	506.6
S.Em\pm	0.29	0.27	0.39	0.56	1.27
LSD @ 5%	0.82	0.78	1.11	1.62	3.62
Pesticides (P)					
P ₀	391.3	410.3	445.6	419.1	359.7
P ₁	402.4	420.7	456.1	441.8	388.7
P ₂	415.3	433.6	470.7	450.7	401.2
P ₃	488.9	506.5	543.0	513.2	447.6
P ₄	458.9	476.9	512.2	493.8	428.9
Mean	431.4	449.6	485.5	463.7	405.2
S.Em\pm	0.26	0.24	0.35	0.50	1.13
LSD @ 5%	0.73	0.70	0.99	1.44	3.24
Interactions (V×P)					
V ₁ P ₀	476.8	511.0	566.8	517.3	401.5
V ₁ P ₁	514.4	546.2	604.2	586.6	496.5
V ₁ P ₂	535.4	567.8	627.8	604.2	516.2
V ₁ P ₃	723.0	753.5	813.5	776.3	655.5
V ₁ P ₄	687.7	717.7	774.6	735.7	633.3
V ₂ P ₀	684.4	716.7	776.7	722.7	645.0
V ₂ P ₁	711.5	741.5	799.6	771.5	683.5
V ₂ P ₂	741.5	771.5	831.5	786.6	720.5
V ₂ P ₃	836.0	866.0	926.0	853.7	770.4
V ₂ P ₄	777.5	807.5	867.5	843.8	739.3
V ₃ P ₀	693.8	723.8	783.8	729.9	645.5
V ₃ P ₁	723.3	753.3	813.3	786.3	698.3
V ₃ P ₂	739.5	769.5	829.5	805.8	717.8
V ₃ P ₃	820.5	850.5	910.5	869.6	778.4
V ₃ P ₄	773.3	803.3	861.7	838.0	750.0
V ₄ P ₀	753.5	783.5	843.5	824.3	705.9
V ₄ P ₁	733.7	763.7	823.7	801.2	713.2
V ₄ P ₂	752.0	782.0	849.0	808.2	720.2
V ₄ P ₃	880.0	906.5	970.0	921.3	779.8
V ₄ P ₄	821.0	851.0	911.0	874.5	736.9
Mean	718.9	749.3	809.2	772.9	675.4
S.Em\pm	2.55	2.43	3.46	5.05	11.32
LSD @ 5%	7.31	6.96	9.91	14.45	32.42

Pesticides were detected only at 24 and 96 hr after the spray. No further detection was seen

V₁ – Malapur local

P₀ – Control (water sprayed)

V₂ – Kalpataru

P₁ – Thiodicarb 75 WP @ 1g/L

V₃ – Manjula

P₂ – Spinosad 45 SC @ 0.1 ml/L

V₄ – Manjari

P₃ – Profenophos 50 EC @ 2 ml/LP₄ – Chlorantraniliprole 20 SC @ 0.5 ml/L

Table 2(a). Influence of pesticides on magnesium content (mg 100 g⁻¹) of brinjal in different growing seasons

Pooled data (kharif 2013-14 & 14-15)					
Treatments	DAT				
	70	80	90	115	125
Varieties(V)					
V ₁	9.08	9.95	10.44	9.46	9.44
V ₂	9.85	9.99	11.28	9.79	9.76
V ₃	9.77	9.99	10.99	9.85	9.87
V ₄	10.45	10.80	11.70	10.65	10.53
Mean	9.79	10.18	11.10	9.94	9.90
S.Em±	0.05	0.01	0.01	0.04	0.03
LSD @ 5%	0.14	0.04	0.03	0.12	0.08
Pesticides(P)					
P ₀	6.73	6.93	7.77	6.82	6.65
P ₁	7.65	7.93	8.62	7.74	7.69
P ₂	7.91	8.32	8.87	8.10	8.08
P ₃	8.64	9.06	9.90	8.77	8.85
P ₄	8.23	8.49	9.25	8.33	8.33
Mean	7.83	8.15	8.88	7.95	7.92
S.Em±	0.04	0.01	0.00	0.04	0.03
LSD @ 5%	0.13	0.03	0.01	0.11	0.07
Interactions(V×P)					
V ₁ P ₀	10.31	11.17	12.24	10.80	10.44
V ₁ P ₁	11.93	13.02	13.75	12.21	12.42
V ₁ P ₂	12.37	13.97	14.12	12.96	13.19
V ₁ P ₃	13.11	14.53	15.24	13.95	13.54
V ₁ P ₄	12.85	13.62	14.26	13.18	13.33
V ₂ P ₀	11.48	11.67	13.44	11.63	11.35
V ₂ P ₁	12.83	12.85	14.45	12.63	12.32
V ₂ P ₂	12.80	13.22	14.91	13.04	13.17
V ₂ P ₃	14.89	15.15	16.88	14.22	14.97
V ₂ P ₄	13.64	13.71	15.53	13.72	13.29
V ₃ P ₀	10.98	11.25	12.84	11.08	11.22
V ₃ P ₁	12.48	13.30	14.22	12.85	13.26
V ₃ P ₂	13.47	13.35	14.48	13.33	12.84
V ₃ P ₃	14.45	14.69	16.24	14.48	14.35
V ₃ P ₄	13.76	14.02	15.50	13.90	14.14
V ₄ P ₀	12.09	12.09	13.32	11.96	11.34
V ₄ P ₁	13.75	13.69	15.02	13.89	13.28
V ₄ P ₂	14.08	14.92	15.65	14.66	14.65
V ₄ P ₃	15.15	16.06	17.65	15.79	16.15
V ₄ P ₄	14.59	15.25	16.36	14.71	14.79
Mean	13.05	13.58	14.80	13.25	13.20
S.Em±	0.44	0.12	0.04	0.39	0.25
LSD @ 5%	NS	0.33	0.10	NS	0.72

Pesticides were detected only at 24 and 96 hr after the spray. No further detection was seen

NS – Non-significant

V₁ – Malapur local

P₀ – Control (water sprayed)

V₂ – Kalpataru

P₁ – Thiodicarb 75 WP @ 1g/L

V₃ – Manjula

P₂ – Spinosad 45 SC @ 0.1 ml/L

V₄ – Manjari

P₃ – Profenophos 50 EC @ 2 ml/LP₄ – Chlorantraniliprole 20 SC @ 0.5 ml/L

Table 2(b). Influence of pesticides on magnesium content (mg 100 g⁻¹) of brinjal in different growing seasons

Pooled data (rabi 2013-14 & 14-15)					
Treatments	DAT				
	70	80	90	115	125
Varieties(V)					
V ₁	9.17	9.96	9.97	9.38	9.49
V ₂	9.90	9.95	10.13	9.89	9.81
V ₃	9.83	9.95	10.02	9.89	9.85
V ₄	10.46	10.76	10.87	10.59	10.56
Mean	9.84	10.16	10.25	9.94	9.93
S.Em±	0.02	0.01	0.01	0.02	0.01
LSD @ 5%	0.07	0.03	0.03	0.06	0.04
Pesticides(P)					
P ₀	6.79	6.93	6.93	6.81	6.67
P ₁	7.66	7.90	8.07	7.72	7.72
P ₂	7.93	8.27	8.39	8.05	8.11
P ₃	8.78	9.06	9.11	8.90	8.88
P ₄	8.20	8.48	8.51	8.26	8.33
Mean	7.87	8.13	8.20	7.95	7.94
S.Em±	0.02	0.01	0.01	0.02	0.01
LSD @ 5%	0.06	0.03	0.03	0.05	0.03
Interactions(V×P)					
V ₁ P ₀	10.57	11.18	11.03	10.62	10.32
V ₁ P ₁	11.70	13.08	13.19	11.82	12.35
V ₁ P ₂	12.36	14.05	13.97	12.74	13.38
V ₁ P ₃	13.52	14.54	14.74	14.14	13.76
V ₁ P ₄	12.98	13.55	13.55	13.21	13.45
V ₂ P ₀	11.40	11.60	11.76	11.45	11.44
V ₂ P ₁	13.04	12.64	13.17	13.00	12.52
V ₂ P ₂	12.83	13.22	13.60	12.99	13.14
V ₂ P ₃	15.15	15.17	15.16	14.88	15.04
V ₂ P ₄	13.57	13.72	13.83	13.60	13.27
V ₃ P ₀	11.10	11.25	11.42	11.20	11.22
V ₃ P ₁	12.43	13.29	13.49	12.65	13.24
V ₃ P ₂	13.48	13.01	13.29	13.39	12.86
V ₃ P ₃	14.59	14.66	14.79	14.66	14.25
V ₃ P ₄	13.91	14.14	13.85	14.04	14.13
V ₄ P ₀	12.17	12.14	11.97	12.12	11.49
V ₄ P ₁	13.91	13.64	13.94	14.03	13.35
V ₄ P ₂	14.18	14.84	15.05	14.56	14.69
V ₄ P ₃	15.26	16.04	16.02	15.67	16.13
V ₄ P ₄	14.21	15.10	15.48	14.19	14.72
Mean	13.12	13.54	13.66	13.25	13.24
S.Em±	0.21	0.10	0.11	0.18	0.11
LSD @ 5%	0.59	0.28	0.30	0.53	0.31

Pesticides were detected only at 24 and 96 hr after the spray. No further detection was seen

V₁ – Malapur local

P₀–Control (water sprayed)

V₂ – Kalpataru

P₁– Thiodicarb 75 WP @ 1g/L

V₃ – Manjula

P₂– Spinosad 45 SC @ 0.1 ml/L

V₄ – Manjari

P₃– Profenophos 50 EC @ 2 ml/L

P₄– Chlorantraniliprole 20 SC @ 0.5 ml/L

Table 3(a). Influence of pesticides on marketable fruit yield (kg plant⁻¹) of brinjal in different growing seasons

Pooled data (kharif 2013-14 & 14-15)						
Treatments	DAT					Total
	70	80	90	115	125	
Varieties (V)						
V ₁	0.55	0.80	0.80	0.60	0.24	2.99
V ₂	0.68	0.90	0.91	0.75	0.43	3.68
V ₃	0.67	0.92	0.82	0.68	0.42	3.52
V ₄	0.75	0.95	0.98	0.87	0.58	4.12
Mean	0.66	0.89	0.88	0.72	0.42	
S.Em±	0.03	0.03	0.01	0.03	0.01	
LSD @ 5%	0.08	0.09	0.03	0.09	0.02	
Pesticides (P)						
P ₀	0.28	0.46	0.37	0.25	0.17	1.54
P ₁	0.54	0.71	0.67	0.52	0.29	2.73
P ₂	0.57	0.76	0.76	0.63	0.37	3.10
P ₃	0.66	0.85	0.93	0.78	0.46	3.68
P ₄	0.60	0.78	0.80	0.71	0.38	3.27
Mean	0.53	0.71	0.70	0.58	0.33	
S.Em±	0.02	0.01	0.02	0.03	0.01	
LSD @ 5%	0.06	0.02	0.07	0.09	0.03	
Interactions (V×P)						
V ₁ P ₀	0.30	0.64	0.43	0.36	0.14	1.88
V ₁ P ₁	0.80	1.04	1.09	0.70	0.24	3.87
V ₁ P ₂	0.84	1.16	1.18	0.83	0.34	4.34
V ₁ P ₃	0.95	1.28	1.43	1.08	0.48	5.23
V ₁ P ₄	0.77	1.20	1.21	1.01	0.42	4.61
V ₂ P ₀	0.56	0.82	0.73	0.46	0.33	2.91
V ₂ P ₁	0.87	1.15	1.15	0.85	0.43	4.46
V ₂ P ₂	0.98	1.28	1.24	1.07	0.70	5.27
V ₂ P ₃	1.12	1.39	1.55	1.36	0.75	6.17
V ₂ P ₄	1.05	1.33	1.42	1.25	0.66	5.72
V ₃ P ₀	0.49	0.79	0.61	0.41	0.27	2.57
V ₃ P ₁	0.91	1.28	1.02	0.84	0.56	4.62
V ₃ P ₂	0.97	1.33	1.18	0.99	0.70	5.17
V ₃ P ₃	1.11	1.46	1.38	1.28	0.82	6.05
V ₃ P ₄	0.99	1.29	1.26	1.03	0.47	5.04
V ₄ P ₀	0.54	0.83	0.67	0.46	0.38	2.87
V ₄ P ₁	1.01	1.29	1.18	1.07	0.70	5.24
V ₄ P ₂	1.03	1.32	1.44	1.32	0.75	5.87
V ₄ P ₃	1.23	1.53	1.80	1.49	1.04	7.09
V ₄ P ₄	1.18	1.37	1.43	1.45	0.99	6.41
Mean	0.89	1.19	1.17	0.97	0.56	
S.Em±	0.02	0.01	0.04	0.03	0.03	
LSD @ 5%	0.07	0.02	0.11	0.09	0.10	

Pesticides were detected only at 24 and 96 hr after the spray.

No further detection was seen

V₁ – Malapur localP₀ – Control (water sprayed)V₂ – KalpataruP₁ – Thiodicarb 75 WP @ 1g/LV₃ – ManjulaP₂ – Spinosad 45 SC @ 0.1 ml/LV₄ – ManjariP₃ – Profenophos 50 EC @ 2 ml/LP₄ – Chlorantraniliprole 20 SC @ 0.5 ml/L

Table 3(b). Influence of pesticides on marketable fruit yield (kg plant⁻¹) of brinjal in different growing seasons

Pooled data (rabi 2013-14 & 14-15)						
Treatments	DAT					
	70	80	90	115	125	Total
Varieties (V)						
V ₁	0.27	0.39	0.61	0.65	0.50	2.42
V ₂	0.46	0.54	0.76	0.75	0.63	3.14
V ₃	0.50	0.52	0.70	0.68	0.57	2.99
V ₄	0.61	0.70	0.83	0.77	0.61	3.53
Mean	0.46	0.54	0.73	0.71	0.58	
S.Em±	0.02	0.01	0.01	0.03	0.01	
LSD @ 5%	0.06	0.02	0.03	0.09	0.03	
Pesticides (P)						
P ₀	0.23	0.20	0.33	0.39	0.32	1.46
P ₁	0.31	0.38	0.55	0.57	0.43	2.23
P ₂	0.37	0.45	0.60	0.60	0.48	2.51
P ₃	0.51	0.61	0.77	0.69	0.57	3.15
P ₄	0.42	0.52	0.65	0.61	0.52	2.72
Mean	0.37	0.43	0.58	0.57	0.46	
S.Em±	0.02	0.03	0.01	0.02	0.02	
LSD @ 5%	0.04	0.09	0.03	0.04	0.06	
Interactions (V×P)						
V ₁ P ₀	0.21	0.24	0.32	0.60	0.28	1.64
V ₁ P ₁	0.27	0.44	0.83	0.87	0.68	3.09
V ₁ P ₂	0.38	0.54	0.89	0.88	0.72	3.41
V ₁ P ₃	0.51	0.76	1.06	1.01	0.83	4.18
V ₁ P ₄	0.44	0.63	0.97	0.95	0.80	3.78
V ₂ P ₀	0.41	0.35	0.59	0.76	0.75	2.86
V ₂ P ₁	0.46	0.59	0.94	0.95	0.74	3.67
V ₂ P ₂	0.63	0.78	0.98	1.05	0.86	4.31
V ₂ P ₃	0.83	0.99	1.39	1.17	1.02	5.39
V ₂ P ₄	0.74	0.90	1.15	1.04	0.88	4.71
V ₃ P ₀	0.47	0.33	0.55	0.44	0.51	2.31
V ₃ P ₁	0.67	0.65	0.87	0.97	0.73	3.89
V ₃ P ₂	0.77	0.80	1.00	1.00	0.81	4.38
V ₃ P ₃	0.89	0.99	1.26	1.13	0.93	5.20
V ₃ P ₄	0.54	0.71	1.01	1.03	0.84	4.13
V ₄ P ₀	0.43	0.39	0.74	0.79	0.61	2.95
V ₄ P ₁	0.70	0.83	1.00	1.00	0.71	4.24
V ₄ P ₂	0.72	0.91	1.16	1.04	0.84	4.66
V ₄ P ₃	1.17	1.35	1.43	1.25	1.01	6.21
V ₄ P ₄	1.08	1.22	1.24	1.05	0.92	5.50
Mean	0.62	0.72	0.97	0.95	0.77	
S.Em±	0.01	0.01	0.02	0.01	0.03	
LSD @ 5%	0.02	0.02	0.05	0.04	0.10	

Pesticides were detected only at 24 and 96 hr after the spray.

No further detection was seen

V₁ – Malapur localP₀ – Control (water sprayed)V₂ – KalpataruP₁ – Thiodicarb 75 WP @ 1g/LV₃ – ManjulaP₂ – Spinosad 45 SC @ 0.1 ml/LV₄ – ManjariP₃ – Profenophos 50 EC @ 2 ml/LP₄ – Chlorantraniliprole 20 SC @ 0.5 ml/L