



SUBLETHAL EFFECT OF RAW AND NEUTRALIZED PICKLING WASTEWATER ON THE HAEMATOLOGY OF *HETEROPNEUSTESFOSSILIS*

Shashikant R. Sitre¹ and Rekha Sarkar²

¹Department of Zoology, N.S.Science and Arts College, Bhadrawati, Dist.Chandrapur .

²CSIR-National Environmental Engineering Research Institute, Nehru Marg, Nagpur.

Corresponding Email: shashikantsitre2008@gmail.com

Communicated : 26.01.2023

Revision : 01.03.2023 & 10.03.2023

Published : 30.05.2023

Accepted : 07.04.2023

ABSTRACT:

Based on recorded LC50 values after experimentation sublethal effects of raw and neutralized pickling wastewater of a steel processing industry of Vidarbha region were analyzed using freshwater catfish *H.fossilis* in long duration experiments for toxicity testing using haematology as a diagnostic tool in national environmental engineering research institute laboratory in nagpur. 1/10th of 96 hrs. LC50 concentration was selected and used for both raw and neutralized pickling wastewater (0.05% in case of raw wastewater and 1.1% in case of neutralized wastewater) for experimentation purpose.

H.fossilis were exposed in a continuous flow (dynamic) bioassay test system and observations on haematological changes in fish blood were recorded after 96 hrs. duration. The general haematological parameters studied include total erythrocyte count, total leucocyte count, haemoglobin content, packed cell volume, clotting time and ESR.

The data on fish haematology was analyzed for statistical significance between the control having normal blood chemistry with fish under treatment in chronic exposures by applying students 't' test and significant differences were recorded at 0.01,0.05 and 0.001 levels between control and treated fish blood.

Keywords :- Sublethal effect, raw pickling wastewater, neutralized pickling wastewater, haematology, *H.fossilis*.

INTRODUCTION :

Human eagerness to perform better and better with respect to production of food, energy and convenience products to enhance the standard of living resulted in increasing industrialization, urbanization and rapid development which in turn has led to tremendous growth in production of various industrial products. Pure chemicals and other toxicants do not enter the aquatic environment as single entities. The concern for environmental protection especially in the aquatic environment arises because industrial effluents contain complex mixtures of several chemicals which are discharged into surface water bodies and subsequently disturb the fragile ecology of receiving systems.

The disposal of industrial wastes in aquatic bodies poses serious problems due to their diverse chemical composition and complexity. The untreated industrial wastewaters often

contain many toxicants which poses threat to aquatic life (David et al, 1988). The entry of these toxicants into aquatic environment triggers a series of events which directly or indirectly affect the aquatic life. The possible effects vary and affect metabolic and physiological functions in organisms or changes in properties of ambient medium that affect the aquatic organisms (Newman *et al*, 1992).

MATERIALS AND METHODS :

Raw and neutralized pickling wastewater from a steel processing industry of Vidarbha region was collected and analyzed for its physico-chemical and heavy metal characteristics in NEERI Laboratory of Nagpur (APHA,1989). The industry manufactures cold rolled steel sheets in wide range of thickness. During cold rolling process hot rolled sheets are treated with hydrochloric acid in a pickling process to remove oxide scales

and to impart clean surface properties to the steel sheets.

After a lot of processing spent hydrochloric acid and rinse water are the two major wastewater sources originating from pickling operation while alkaline wastewater stream originates from alkali cleaning. These are the wastewater sources which are collected and analyzed as raw and neutralized by lime. The characterization of raw and neutralized pickling wastewater was undertaken with respect to PH, conductivity, acidity, chlorides, TSS, TDS, oil and Grease & heavy metals iron, zinc, cadmium, chromium, nickel, manganese, lead, aluminium, copper, and silicon. The color of raw pickling wastewater was yellow and that of neutralized wastewater was pale greenish.

Freshwater catfish *H.fossilis* is used for studying the sublethal effect of raw and neutralized pickling wastewater. The fish size varies from 12-17 cms. While weight ranges from 50-60gms. Acute toxicity tests were performed for estimating LC50 values for raw and Neutralized wastewater. The concentration of 1/10th of 96 Hrs LC50 was chosen for raw and neutralized pickling wastewater (Sprague, 1969; APHA,1989; Mckim *et al*,1970). In case of raw pickling wastewater a sub-lethal dose of 0.055 was selected and in case of neutralized wastewater a sub-lethal dose of 1.1% v/v was used for experimentation. The exposed catfish was taken out after 96 hrs. interval and blood samples were obtained and analyzed (Wintrobe,1973; Dacie and Lewis, 1975; Sood,1996).

The haematological parameters investigated include total RBC count, total leucocyte count, haemoglobin content, packed cell volume, clotting time, and ESR. The data was analyzed using students 't' test and significant differences established at 0.01,0.05 and 0.001 levels of significance.

RESULT AND DISCUSSION :

Today no instrumental or chemical analysis is available to test the toxicity of pure chemical or complex wastewater to living organisms. The only real measure of toxicity is through test species whose dose response relationship can be obtained through time, either experimentally or through continuous field monitoring. The results of these toxicity tests are used for establishing, predicting and deriving the maximum permissible concentrations at which the toxicants may be present in the environment without causing damage to the aquatic organisms (Sarkar *et al*,1995).

Sublethal dose of raw (0.05%)v/v and neutralized (1.1%) pickling industry wastewaters produced marked changes in the haematological parameters of *H.fossilis* during 96 hours duration. The changes in raw wastewater are shown in Table 1 and in case of neutralized it was shown in Table2. The impact of the pickling wastewater in case of general haematological parameters was expressed as significant decrease in total erythrocyte count, haemoglobin content, packed cell volume and clotting time with a significant rise in total leucocyte count and erythrocyte sedimentation rate. The degree of impact was found to be considerably reduced on neutralization of raw wastewater as evident from the observations. The Mean Corpuscular Volume (MCV), Mean Corpuscular Haemoglobin (MCH) and Mean Corpuscular Haemoglobin Concentration (MCHC) were calculated with a view to designate the type of anaemia in *H.fossilis* during prolonged exposure to raw and neutralized wastewater.

Blood is an indicator of most of the vital physiological and biochemical functions. Many investigators have emphasized the need for the establishment of normal haematological values in fish from unpolluted environments with a view to diagnose impacts of environmental

stress due to pollution (Larsson *et al*, 1976; Thakur and Sahai,1986; Helawell, 1986).

An inherent need to develop rapid quantitative physiological methods to measure the impact of pollutants present at subacutelevels in the aquatic environment has been expressed by these authors. The leucocytes are involved in the immune defence of an organism and any stress or damage to internal organs, parasitic infection, necrosis, injury or inflammation stimulates the production of leucocytes.

REFERENCES:

- APHA (1989). Standard Methods for the Examination of Water and Wastewater, 17th Edition, American Public Health Association, New York.
- Dacie J.V.and Lewis S.M. (1975). Practical Haematology, Churchill Livingstone, London,UK.
- Helawell J.M.(1986). Biological Indicators of Freshwater Pollution and Environmental Management, Elsevier Applied Science Publishers, London and New York.
- Larsson A, Bengtsson B.E. and Swanberg, O.(1976). Some haematological and biological effects of cadmium on fish In: Effects of pollutants on Aquatic Organisms (Ed. Lockwood A.P.M.) Cambridge University press, London, 2; 34-35.
- Mckim J.M, Christensen G.M. and Hunt E.P.(1970). Changes in the blood of brook trout (*Salvelinus fontinalis*) after short term and long term exposure to copper. J.Fish.Res.Board of Can. 27: 1883-1889.
- Sprague J.B.(1969). Measurement of pollutant toxicity to fish I: Bioassay Methods for Acute toxicity, Water Res. 3 : 793-821.
- Sood R (1996). Haematology for Students and Practitioners, 4th Edition, Jaypee Brothers Pvt. Ltd. New Delhi, pp. 397.
- Thakur N and Sahai S.(1986). Carbaryl induced haematological alterations in the teleost *Garra gotyla gotyla* (Gray) In Proceedings of Symposium on “ Man Development Bioresources and Enviornment”. H.S.Gour Vishwavidyalaya,Sagar,M.P. Dec. 26-28 pp. 339-344.
- Wintrobe, M.M.(1973). Clinical Haematology, 7th Edition, Lee and Fabiger, Philadelphia, pp. 935.
- Sarkar R,Chaudhari P.R.,Sitre S and Gajghate D.G.(1995). Toxicity testing through Fish Bioassay and Its Application in India, in “ Pollution and Biomonitoring “ Tata McGraw Hills Publ.Company, New Delhi.

Table 1 : Alterations in Haematological Parameters of *H.fossilis* during 96 Hrs. exposure to Raw Pickling Wastewater at sublethal dose of 0.05%v/v

Sr.No.	Parameter	Observations		% Changes
		Control	Exposed	Observed
1.	Total erythrocyte Count ($\times 10^6/\text{cm}$)	3.35 \pm 0.05	2.52 \pm 0.04	24.77(-)
2.	Total leucocyte Count ($\times 10^4/\text{cm}$)	4.50 \pm 0.13	6.80 \pm 0.04	51.11(+)
3.	Haemoglobin Content (Gm/100ml)	11.80 \pm 0.05	10.00 \pm 0.11	15.25(-)
4.	Packed Cell Volume (%)	41.00 \pm 0.42	38.50 \pm 0.23	6.09(-)
5.	Clotting Time (sec.)	114.00 \pm 2.07	107 \pm 1.14	6.14(-)
6.	Erythrocyte Sedimentation Rate (mm/Hr)	1.30 \pm 0.11	1.70 \pm 0.08 ²	30.76(+)
7.	Mean Corpuscular Volume (cm^3/cell)	122.38 \pm 1.90	152.77 \pm 1.62 ¹	24.83(+)
8.	Mean Corpuscular Haemoglobin (pg/cell)	35.22 \pm 0.93	39.68 \pm 0.5 ¹	12.66(+)
9.	Mean Corupscular Haemoglobin Concentration (%)	28.78 \pm 0.32	25.97 \pm 0.14 ¹	9.76 (-)

Results are expressed as Mean \pm S.D.

1. P < 0.01
2. P < 0.05

Table 2 : Alterations in Haematological Parameters of *H.fossilis* during 96 Hrs. exposure to Neutralized Pickling Wastewater at sublethal dose of 1.1% v/v

Sr. No.	Parameter	Observations		% Changes
		Control	Exposed	Observed
1.	Total erythrocyte Count ($\times 10^6/\text{cm}$)	3.65 \pm 0.10	2.79 \pm 0.07	23.56(-)
2.	Total leucocyte Count ($\times 10^4/\text{cm}$)	4.20 \pm 0.16	6.20 \pm 0.08	47.61(-)
3.	Haemoglobin Content (Gm/100ml)	11.70 \pm 0.07	10.10 \pm 0.08	13.879(-)
4.	Packed Cell Volume (%)	40.50 \pm 0.21	38.90 \pm 0.73	4.96 (-)
5.	Clotting Time (sec.)	111.00 \pm 1.58	105.10 \pm 1.30	5.10(-)
6.	Erythrocyte Sedimentation Rate (mm/Hr)	1.20 \pm 0.13	1.50 \pm 0.10	25.00(-)
7.	Mean Corpuscular Volume (cm^3/cell)	110.00 \pm 1.20	139.42 \pm 0.78	26.74(+)
8.	Mean Corpuscular Haemoglobin (pg/cell)	32.05 \pm 0.52	36.20 \pm 1.45	12.94(+)
9.	Mean Corupscular Haemoglobin Concentration (%)	28.88 \pm 0.31	25.96 \pm 0.15	10.11(-)

Results are expressed as Mean \pm S.D.

1. P < 0.01
2. P < 0.05