



## Ultrasonic Studies of The System Containing Aqueous Solution of Sodium Salt of 4-Amino Salicylic Acid At Different Temperatures

Pravin J. Ganjare<sup>1</sup> Shashikant R. Aswale<sup>2</sup> and Sunanda S. Aswale<sup>3</sup>

<sup>1</sup>Shivramji Moghe Mahavidyalaya, Pandharkawada, Dist. Yavatmal, M. S

<sup>2</sup>Sant Gadge Baba Amravati University, Amravati, Dist. Amravati., M.S.

<sup>3</sup>Lokmanya Tilak Mahavidyalaya, Wani. Dist. Yavatmal, M. S. 445304.

Email-[ganjarepravin@yahoo.com](mailto:ganjarepravin@yahoo.com), [ssaswale@rediffmail.com](mailto:ssaswale@rediffmail.com)., Mob.09673560433.

### Abstract:-


When acoustic waves are propagated through a liquid, dissipation of energy in sound waves takes place, the study of molecular interactions in various organic liquids and liquid mixtures is of considerable importance in recent years. 4-amino salicylic acid is most commonly used in the treatment of tuberculosis, known as Para- Amino Salicylic acid (PAS). It is antibiotic drug always used as Pro-drug. It is the most important member in the treatment of multidrug resistant treatment with few side effects. In the present work, ultrasonic velocity, density and viscosity of aqueous solutions of sodium salt of 4-amino salicylic acid have been measured experimentally using ultrasonic interferometer with 4 MHz frequency over entire range of concentration at temperatures 298K, 303K and 308K. From this experimental data, other physical parameters such as acoustical impedance (Z), adiabatic compressibility ( $\beta$ ), intermolecular free length (Lf), free volume (Vf), Rao's constant (R), Wada's constant (W) have been determined for each composition. The computed acoustical parameters and their values point to the presence of specific molecular interaction in the mixtures and has been interpreted in light of solute-solvent interactions. Hence it is concluded that the association in these mixtures is the result of Hydrogen bonding in Binary liquid mixtures. The results may be helpful to understand the exact mode of action of the drug in vivo.

**Keywords:-** Ultrasonic velocity, free volume (Vf), Rao's constant (R), Wada's constant (W).

### Introduction:-

The study of intermolecular interactions plays an important role in the development of molecular sciences. A large number of studies have been made on the molecular interaction in liquid system by various physical methods like Infrared<sup>1,2</sup>, Raman effect<sup>3,4</sup>, Nuclear magnetic resonance, Dielectric constant<sup>5</sup>, Ultraviolet spectroscopy<sup>6</sup> and Ultrasonic method<sup>7,8</sup>. In recent years, Ultrasonic technique has become a powerful tool in providing information regarding the molecular behavior of the medium. Ultrasonic's is an area of intense scientific and technological research. Science and technology of ultrasonic is widely sought in the recent years for industrial and medical application. Research and development in ultrasonic's have also maintained a steady pace for the past several decades to meet intense demands. The change in the wavelength of ultrasound waves in different media is due to the elastic properties and the induced particles vibrations in the medium. Further the wavelength of ultrasonic wave is small and hence it exhibits some unique phenomenon in addition to the properties of sound waves. The study of propagation of ultrasonic wave in liquid systems and solid is now rather well established. The ultrasonic wave is an effective means for examining certain physical properties of the material. The ultrasonic velocity and density measurement of certain dielectric liquids can be used to compute acoustical parameters such as

adiabatic compressibility, free length, Acoustic impedance, Rao's Constant, and Wada's constant<sup>9,10</sup>. The present investigation deals with the study of molecular interaction in Sodium 4-Amino-Salicylate dihydrate in water at 298K, 303K and 308 K. further the data has been obtained at different concentration with a view to understand the effect of temperature on these properties



### Sodium 4-Amino Salicylate di hydrate

#### Experimental Section:-

##### Materials:-

Analytical Range (AR) Sodium 4-Amino Salicylate dihydrate is used in the present work. The solutions were prepared by using double distilled water as solvent. The concentration range selected was 0.1M, 0.01M and 0.001M.

##### Methods:-

All the weighings was done on digital electronic balance Model-CB/CA/CT-Series Contech having accuracy  $\pm 0.0001$ g.

The densities of the solutions were measured accurately using digital densitometer (Model-DMA Anton Paar). Viscosity of the solutions was measured by Ostwald’s viscometer which was calibrated with benzene and double distilled water at all three temperatures. The values are accurate to ± 0.001 cp.

The ultrasonic velocity was measured by using ultrasonic multi frequency interferometer (Model- M-83) supplied by Mittal Enterprises New Delhi, operating at 4MHz frequency with an accuracy of ± 2 m/s. The principle used in the measurement of ultrasonic velocity through medium is based on the accurate determination of wavelength of ultrasonic waves of known

frequency produced by quartz crystal in measuring cell. The temperature of the solution was maintained by circulating water through the jacket of double walled cell. Measurements were made using constant temperature bath within ± 0.01K.

**RESULTS AND DISCUSSION:**

The experimentally measured values of density, viscosity and sound speed of solutions of 4 - amino salicylic acid sodium salt at 298K, 303K and 308K are given in **Table 1**.

The acoustical parameters were calculated from v, η and ρ values using standard formulae, and given in **Table 2,3 and 4**.

1) Adiabatic Compressibility -  $\beta = 1/v^2 \rho$  ...**(1)**

2) Intermolecular free length -  $L_f = K \sqrt{\beta_s}$  ...**(2)**

3) Specific acoustic impedance -  $Z = v_s \cdot \rho$  ...**(3)**

4) Rao’s Constant -  $R = (M_{eff} / \rho) \times v^{1/3}$  ...**(4)**

5) Wada’s Constant -  $W = (M_{eff} / \rho) \times \beta^{-1/7}$  ...**(5)**

6) Relative Association -  $RA = \rho_s / \rho_o [v_o / v_s]^{1/3}$  ...**(7)**

7) Relaxation time -  $\tau = 4/3 \beta \times \eta$  ...**(8)**

8) Free Volume -  $V_f = M_{eff} \times v_s / k \times \eta$  ....**(9)**

Where  $k = 4.28 \times 10^9$ , Temperature Independent Constant for all liquids.

9) Molar Conductance -  $\mu_v = Kc[1000/M]$  ....**(10)**

**Table 1:** Density, Velocity and Viscosity at 298, 303, 308 K (At Frequency-4MHz)

Sr. No.	Temperature (K.)	Concentration(M)	Density(ρs) (Kg/m <sup>3</sup> )	Velocity(v <sub>s</sub> )(m/s)	Viscosity(η) (Pa.S.) or Kg m <sup>-1</sup> s <sup>-1</sup>
1	298	0.1	1005	9105.44	8.93E-04
2		0.01	998.6	9991.6	8.64E-04
3		0.001	997.4	10812.23	8.39E-04
4	303	0.1	1004.7	9444.4	8.46E-04
5		0.01	998.3	10500.5	8.02E-04
6		0.001	997.01	11312.26	7.70E-04
7	308	0.1	1003.5	9754.9	7.42E-04
8		0.01	997.7	11091.16	7.16E-04
9		0.001	996.6	12015.26	6.64E-04

**Table 2 :-** Adiabatic Compressibility, Acoustic impedance and Free length at .298, 303, 3080 K.


Sr.No.	Temperature (K)	Concentration M	Adiabatic Compressibility (βs) Pa-1	Acoustic Impedance Z x(Kgm <sup>-2</sup> S <sup>-1</sup> )	Free length L <sub>f</sub> (m)
1	298	0.1	1.2E-11	9150967	6.81E-12
2		0.01	1.00E-11	9977612	6.23E-12
3		0.001	8.58E-12	10784118	5.76E-12
4	303	0.1	1.12E-11	9488789	6.61E-12
5		0.01	9.08E-12	10482649	5.97E-12
6		0.001	7.84E-12	11278436	5.54E-12
7	308	0.1	1.05E-11	9789042	6.48E-12
8		0.01	8.15E-12	11065650	5.71E-12
9		0.001	6.95E-12	11974408	5.28E-12

**Table 3 :** Relative association, and relaxation time


Sr. No.	Temperature(K)	Concentration(M)	Relative Association (R <sub>A</sub> )	Relaxation Time τ
1	298	0.1	8.45E-01	1.43E-14
2		0.01	0.83926	1.16E-14
3		0.001	0.791	9.59E-15
4	303	0.1	0.85143	1.26E-14
5		0.01	0.813319	9.71E-15
6		0.001	0.793701	8.04E-15
7	308	0.1	0.859373	1.04E-14
8		0.01	0.815329	7.78E-15
9		0.001	0.793701	6.15E-15

**Table 4 :** Rao's Constant, Wada's Constant, Free volume and Molar conductance.


Sr.No.	Temperature(K)	Concentration(M)	Rao's Constant (R)	Wada's Constant (W)	Free Volume (V <sub>f</sub> )	μ=Kc*(1000/M)mhos mol.
1	298	0.1	2.08E-03	3.61E-03	3.67E-06	5.10E+04
2		0.01	0.00215	3.72E-03	4.43E-06	1.06E+05
3		0.001	2.21E-03	3.81E-03	5.20E-06	2.04E+05
4	303	0.1	2.10E-03	3.65E-03	4.20E-06	5.51E+04
5		0.01	2.19E-03	3.77E-03	5.33E-06	1.33E+05
6		0.001	2.24E-03	3.86E-03	6.33E-06	2.60E+03
7	308	0.1	2.13E-03	3.68E-03	5.37E-06	6.28E+04
8		0.01	2.23E-03	3.83E-03	6.85E-06	1.52E+05
9		0.001	2.29E-03	3.93E-03	8.65E-06	3.50E+04




**Figure -1** Velocity vs conc.



**Figure -2** Adiabatic compressibility vs conc.



**Figure -3** Acoustic Impedance vs conc.



**Figure -4** Free length vs conc.




Figure -5 Relative association vs conc.




Figure -6 Relaxation time vs conc.




Figure -7 Rao's Constant vs conc.




Figure -8 Wada's Constant vs conc.




Figure -9 Free volume vs conc.




Figure -10 Molar conductance vs conc.

This paper deals with the study of molecular interaction in binary liquid mixtures of Sodium 4-amino salicylate dihydrate at 298K, 303K and 308K. We have reported ultrasound velocity ( $V_s$ ), density ( $\rho_s$ ) and viscosity ( $\eta$ ) of binary liquid mixture. With the help of experimental data, the thermodynamic and acoustic properties like adiabatic compressibility ( $\beta_s$ ), intermolecular free length ( $L_i$ ), free volume ( $V_i$ ), and their excess values have been computed at three different temperatures 298K, 303K, 308K. The above functions and their excess values are tabulated and graphed in figures. The 4-amino salicylate in water acts as polar molecule.

The adiabatic compressibility is the fractional decrease of volume per unit increase of pressure, when no heat flows in or out. These changes are related to the compressibility of the medium by thermodynamic relation

$$\beta = 1/v^2 \rho_s$$

From the **Table.1** it is noted that the density and viscosity decreases with decrease in concentration for all the cases. Ultrasonic velocity decreases with increase in concentration (**Fig.1**) of the solute in all the systems. The decrease in velocity may be due to the increase in free length and adiabatic compressibility of the liquid mixtures (**Fig.2 and 4**) which shows the change in the structural arrangement of the molecules in the mixture as stated in some other aqueous binary mixtures<sup>11,12</sup>. The adiabatic compressibility and free length increases with increase in concentration for all the systems. This may lead to the presence of specific molecular interaction between the molecules of the liquid mixture. The adiabatic compressibility and free length are the deciding factors of the ultrasonic velocity in liquid systems.

According to Eyrings liquid state theory, the acoustic wave, which was excited in the liquid, is transmitted momentarily to the intermolecular free length and the rate process in liquids is determined by the free volume. So free volume plays an important role in ultrasound wave propagation. The values are reported in Table 4. Free volume decreases with increasing concentration at constant temperature.

The decrease in free volume shows that the strength of interaction decreases gradually with the increase in solute concentration. It represents that there is weak interaction between the solute and solvent molecules. Acoustic impedance decreases with increase of concentration for all the three systems. **(Fig. 3)** The relaxation time ( $\tau$ ) increases with increasing concentration for all the three systems. **(Fig. 6)** The dispersion of the ultrasonic velocity in the system should contain information about the characteristic time of the relaxation process that causes dispersion. The relaxation time which is in the order of  $10^{-14}$  sec is due to structural relaxation process<sup>13</sup> and in such a situation it is suggested that the molecules get rearranged due to co-operative process<sup>14</sup>.

Rao's Constant and Wada's Constant decreases with increase in concentration at all temperatures. The variation of molar sound velocity or Rao's Constant and molar compressibility or Wada's Constant are found to be linear **(Table.4 and Fig. 7 and 8)**. It shows the presence of solute-solvent interaction.<sup>15</sup>

The relative association depends on either the breaking up of the solvent molecules on addition of solute to it or the salvation of ions that are present. In the present case relative association increases with increase in concentration **(Fig.5)** which indicates prominent solute-solvent interaction.

#### Conclusion:-

The computed acoustical parameters and their values point to the presence of specific molecular interaction in the mixtures. Hence it is concluded that the association in these mixtures is the result of Hydrogen bonding in Binary liquid mixtures.

#### References:-

- [1]. Eyring, H. and Kincaid, J.F., J. Chem. Phys., 6,620.(1938).
- [2]. Singh, S., Singh, R., Prasad, N. and Prakash, S., Ind. J. Pure and Appl. Phys., 3,156 (1977).
- [3]. Ramamurthy, M. and Sastry, O.S., Indian J. Pure and Appl. Phys., 21, 579, (1983).
- [4]. (i) Fletcher, A., J. Phys. Chem., 73, 2217, (1969).  
(ii) Hamaker, R., Clegg, R., Patterson, P., Riddick, P. and Rock, P., and Rock, S., J. Phys. Chem., 72,1837, (1968).  
(iii) Ramasamy, K. and Ranganathan, V., Indian J. Pure and Appl. Phys., 8,144, (1970).
- (iv) Venkateswaran, K., Krishnapillai, M.G. and Ramasamy, K., Proc. Indian Acad. Sci., 53,195, (1961).
- [5]. Hobbs, M.E. and Bates, W.W., J. Am. Chem. Soc., 74,746, (1952).
- [6]. Negakuva., J. Am. Chem. Soc., 76,3070, (1954).
- [7]. Freedman, E., J. Chem. Phys., 21, 1784, (1955).
- [8]. Kannappan, A.N. and Rajendran, V., Indian J. Pure and Appl. Phys., 30,176, (1992).
- [9]. Aswale S.R., S.S. Aswale and R.S. Hajare., Int. J. Pharm. Pharm. Sci., Vol.5, Suppl.1, 76-79 (2013)
- [10]. Aswale S.S., Aswale S.R. and Dhote A.B.- Int. J. Res. Chem. Environ., Vol.2, issue 4 Oct. 2012 (154-158)
- [11]. Syal, V.K., Anitha Chawla and Suvarcha Chawla, J. Pure Appl. Ultrason. 27 (2005) 61-69
- [12]. Kothai, S. and Kannapan, V., Ind. J. Pure and Appl. Phys. 40 (2002) 17.
- [13]. Kinsler, L.E. and Ray, A.R., Fundamentals of Acoustics (Wiley Eastern, New Delhi) (1989).
- [14]. Ali, A., Hyder, S. and Nain, A.K., Ind. J. Phys. 74 B, 63 (2000).
- [15]. Jaykumar, S., et al., J. Acou. Soc. India, 28, 1-4; 373-375, 2000.