



MEASUREMENT OF GROWTH AND PHYCOBILINS IN A CYANOBACTERIUM *PHORMIDIUM FRAGILE* (MENEHINI) GOMONT

Janardhan Namdeo Nehul

Department of Botany, Dada Patil Rajale College, Adinathnagar Tal-Pathardi Dist-
Ahmednagar- 414505(MS), India

Email: jnnehul@gmail.com

Communicated: 5.03.21

Revision :26.03.21 & 17.04.2021
Accepted: 15.05.2021

Published: 30.05.2021

ABSTRACT:

Phormidium fragile was isolated from the collected soil samples from different locations of Ahmednagar district of Maharashtra state (India). Identification was carried out using morphological variation and taxonomical approaches according to Desikachary. The axenic culture of *Phormidium fragile* was obtained in the laboratory. For the biomass production, different culture media were used namely BG-11, Fogg's medium, Allen and Arnon medium, Zarrouk's medium and CFTRI medium. The biomass was harvested by filtration through double layered muslin cloth and dried using air blower. After harvesting, the biomass obtained was subjected to the growth analysis. Phycobilins were estimated by following the method described by Bennett and Bogorad. Out of the different culture media used, BG-11 medium supported the growth of *Phormidium fragile* properly as compared to other media used. Phycobilins content was found to be more in *Phormidium fragile* grown in BG-11 medium followed by Fogg's medium.

Keywords: *Phormidium fragile*, Phycobilins, BG-11, Fogg's medium, Allen and Arnon medium, Zarrouk's medium and CFTRI medium

INTRODUCTION:

Cyanobacteria (blue-green algae, BGA) are morphologically diverse group of phototrophic prokaryotes, which occur in almost every habitat on earth and useful to mankind in various ways ¹. They constitute a vast potential resource in varied applications such as food, feed, fuel, fertilizer, medicine, industry and in combating pollution ¹. Until past few decades of research, cyanobacteria were of academic interests and were mostly ignored as nuisance but, now are proved as potential organisms for much biotechnological utilization ^{1,2-3}. The interest in these organisms as generators of pharmacologically active and industrially important compounds has been stimulated by recent results ⁴. The carbohydrates produced by

cyanobacteria have important commercial uses. Since carbohydrates are non-toxic, they are desirable and used in the food industry ⁵. Carbohydrates are frequently used in dietary additives for poultry and aquaculture farming ⁶. A large number of marine nitrogen-fixing cyanobacteria have been tested for their nutritional value with the hybrid *Tilapia* fish fry and found that the marine cyanobacterium *Phormidium valderianum* BDU 30501 has shown to serve as a complete aquaculture feed source, based on the nutritional qualities and non-toxic nature with animal model experiments^{1,7}. Several micro algae such as *Chlorella*, *Scenedesmus* and *Coelastrum* have been established as good quality protein sources ⁸. The main advantage of these species

is their high protein content; therefore they are used as food supplements. They also present great benefits to human health due to their antioxidant properties, their role as activator of cell regeneration, and their positive effect on kidney and memory problems ⁹.

Cyanobacteria possess all the known phycobilins such as phycocyanin, phycoerythrin, Phycoerythrocyanin and allo-phycocyanin. Among them, Phycocyanin and phycoerythrin are commercially valuable. Linablu, a phycocyanin product from Dainippon Ink and chemicals Inc., Japan is an odorless, non-toxic blue powder and used for coloring candy, ice-cream, dairy products and soft drinks¹⁰. Phycoerythrin from *Spirulina* and other cyanobacteria is used as a food colour for products like ice-cream ¹¹.

Apart from the use of phycobilins as food grade dyes, they are also used as tools for basic research and medical diagnostics. They are used in fluorescence microscopy and fluorescence immunoassays ¹²⁻¹³. Phycocyanin, the major phycobiliprotein also exhibited anti-cancer activity, stimulation of immune system and ability to treat ulcers and haemorrhoidal bleeding ². The present study was carried out for the estimation of growth and phycobilins content in *Phormidium fragile* in different culture media.

MATERIALS AND METHOD:

Method of collection-The soil samples from 5-10 cm deep soil layers were collected using the scalpels. Soil samples were collected in polythene bags of size 6 x 4 inches.

Nutrient media-The different culture media namely BG-11 ¹⁴; Fogg's medium ¹⁵⁻¹⁶; Allen and Arnon's medium ¹⁷; CFTRI medium ¹⁸ and Zarrouk's medium ¹⁹ were used for the rich growth of *Phormidium fragile*. These media were separately used in different sets.

Isolation of cyanobacterial species-The dry soil samples were spread in petri dishes and moistened with sterilized distilled water and cultures were incubated in light. When the visible growth of cyanobacteria begins to appear in the cultures, these cultures were used for the isolation of unialgal cultures of *Phormidium fragile*.

Identification of the algal samples - Morphometric studies were carried out by using ocular and stage micrometer. The identification of *Phormidium fragile* was carried out using monograph and keys of Desikachary ²⁰.

Biomass production-For production of biomass, glass bottles (300 mL capacity) were used. The bottles were filled with 100 mL medium and autoclaved. The inoculum was ground in the sterile mortar and pestle in laminar air flow. Then the bottles were inoculated with 5 mL of unialgal suspension of *Phormidium fragile* and labeled properly. All the cultures were maintained in the culture room at temperature 28±2°C under 8-h light/16-h dark photoperiod with a photosynthetic photon flux density of 40 µmoles⁻²S⁻¹ provided by cool white fluorescent tube lights. After harvesting, the biomass obtained was subjected to the growth and proteins analysis.

Estimation of phycobilins

Phycobilins were estimated by following the method described by Bennett and Bogorad (1973). Two mL cell suspension was centrifuged at 4500 rpm for 10 minutes and pellet of algal biomass was used for the estimation of total phycobilins. The pellet of biomass of *Phormidium fragile* was blotted with filter papers to remove maximum quantity of water from it. The pellet thus dried was homogenized in 100 µl glycerol and placed in the dark for 3 hours. After the contents were mixed, 5 mL of 10 % ammonium sulphate containing 1mL of 3

mM sodium azide and 1mL of 10 mM Na₂EDTA²¹. The resultant cell suspension was sonicated for 10 minutes and then centrifuged at 4500 rpm for 10 minutes. Absorbance of the supernatant solution was determined at 565 nm, 620 nm and 650 nm on UV Visible spectrophotometer (Systronics, India; model 2202). The amounts of phycoerythrin, phycocyanin and allophycocyanin were calculated by using following equations²².

A. Amount of Phycoerythrin (PE)

$$= A_{565} \cdot 2.8 [PC]^{-1.34} [APC] \text{ mg ml}^{-1} / 12.7$$

B. Amount of Phycocyanin (PC),

$$= A_{620} - 0.7 \times A_{650} \text{ mg ml}^{-1} / 7.38$$

C. Amount of allophycocyanin (APC)

$$= A_{650} \cdot 0.19 \times A_{620} \text{ mg ml}^{-1} / 5.65$$

Where, A_{565} = Absorbance of supernatant solution at 565nm,

A_{620} = Absorbance of supernatant solution at 620 nm,

A_{650} = Absorbance of supernatant solution at 650nm.

D. Sum of PE, PC and APC gives content of total phycobilins.

Amount of Phycobilins are expressed as % of dry weight basis.

RESULTS AND DISCUSSION:

Out of the different culture media used, BG-11 medium supported the growth of *Phormidium fragile* properly as compared to other media used. Allen and Arnon medium also supported growth but after 20 to 25 days, photo bleaching of biomass was observed. Other growth media, such as Fogg's medium and Zarrouk's medium supported the growth of *Phormidium fragile* but the growth rate was very slow.

Yield of biomass is one of the direct measures of quantity of biomass produced per unit area within a specific time. Higher yield indicates higher biomass produced per unit area. Comparison of *Phormidium fragile* in different

media showed that highest biomass per bottle in terms of dry weight was produced in BG-11 medium followed by Allen and Arnon medium. The phycobilins content was found to be more in the *Phormidium fragile* grown in BG-11 medium followed by the Fogg's medium. CFTRI and Zarrouk's medium showed poor response for the phycobilins content as compared to other media.

Values are mean \pm SE of three independent experiments.

Cyanobacteria are photoautotrophic bacteria and require all the essential major and minor elements. The heterocystous cyanobacteria fix atmospheric nitrogen and they can use atmospheric nitrogen as a source of nitrogen. In bottles, the medium does not come in contact with atmospheric nitrogen and the source needs to be added in the culture medium. If the culture medium is devoid of nitrogen, it results in poor growth of cyanobacteria. Similar results were reported by Olatz²³. Medium lacking nitrogen source, results in yellowish green color of the cells which is a characteristic of nitrogen deficiency. In the culture methods like photo- bioreactors, pure nitrogen is continuously bubbled into culture medium²⁴⁻²⁶ so that cultures do not get affected due to nitrogen deficiency.

The growth of *Phormidium fragile* was more in BG-11 medium than in other media. For optimum growth of cyanobacteria, appropriate $Ka^+ : Na^+$ ratio is required in the cytoplasm. Adequate Na^+ is required by nitrogen fixing cyanobacteria for conversion of molecular nitrogen into ammonia²⁷. BG-11 medium consists moderate concentration of Na^+ and in Allen and Arnon medium, Zarrouk's medium and CFTRI medium, there is high concentration of Na^+ while in Fogg's medium;

there is no Na⁺ source. *Phormidium fragile* is from moist soil habitat, which may not require high concentration of Na⁺ ions in the medium.

Influence of culture media on phycobilins Phycoerythrin

The percentage of phycoerythrin was affected by the composition culture media. In all the media used, phycoerythrin was found to be highest in the Fogg's medium in *Phormidium fragile*. The highest amount of phycoerythrin 1.43% was produced in *Phormidium fragile* cultured in BG-11 medium. CFTRI medium supported poorly for the production of phycoerythrin as compared to other media used.

Phycocyanin

The amount of phycocyanin varied from medium to medium. BG-11 and Fogg's medium were the best media for the accumulation of the phycocyanin content in *Phormidium fragile* species. The highest phycocyanin content was found in *Phormidium fragile* cultured in BG-11 medium and lowest 1.22 % in *Phormidium fragile* grown in CFTRI medium. The quantity of phycocyanin produced in BG-11 was found 2.97%. In Fogg's medium the percentage was 1.44 %. There was a production of less amount of phycocyanin in other culture media.

Allophycocyanin

In *Phormidium fragile*, maximum allophycocyanin content was found in the Fogg's medium followed by the BG-11 medium. Other media showed poor response for the allophycocyanin content in *Phormidium fragile*. The maximum quantity 2.58% was occurred in *Phormidium fragile* grown in BG-11 and 2.29% cultured in Fogg's medium. There was an accumulation of less amount of allophycocyanin in *Phormidium fragile* species grown in CFTRI medium.

BG-11 medium was superior for the total phycobilins content in *Phormidium fragile*

species studied followed by the Fogg's medium. CFTRI medium shows very poor response for the total phycobilins content in *Phormidium fragile* species studied. Composition and pH of the medium also affect the content of phycobilins and there is change in the colour of culture of *Phormidium fragile*. The production of phycobilins is greatly affected by the pH. This indicates the inability of the cyanobacteria to maintain a constant internal pH²⁸.

CONCLUSION:

There is an effect of different culture media on growth. It may be because of either composition of culture medium or pH value. The percentage of phycobilins also changed in various culture media used. BG-11 culture medium is suitable for the proper growth and commercial production of phycobilins pigments in a cyanobacterium *Phormidium fragile*. Fogg's medium can be efficiently used for the production of phycobilins pigments used for various purposes.

REFERENCES:

- Thajuddin N, Subramanian G (2005) Cyanobacterial biodiversity and potential applications in biotechnology, Curr. Sci. 89:Pp 47-57
- Richmond A (1990) In: *Algal Biotechnology*, (Eds.). Round FE, Chapman DJ, Bristol UK: Biopress Ltd. Pp. 269-330
- Sundararaman M, Sekar S (2001) Biotechnological potential of cyanobacteria. In: *Algal biotechnology*, Trivedi PC (ed.), Pointer Publishers, Jaipur, India. Pp 31-52
- Singh DP, Tyagi MB, Kumar A, Prasuna EG (2002) Bioactive Secondary Metabolites from cyanobacteria. Pg.275-292. In *Algological research in India*. Edt. N. Anand, Pbl. Bsmpps Dehradun. India

- Bauernfeind JC (1981) Carotenoids as colorants and Vitamins A Precursors. London: Academic Press
- Hirschberg J, Chamoritz D (1994) In: *The Molecular Biology of Cyanobacteria*, (Ed.). Bryant, DA Dordrecht, The Netherlands: Kluwer Academic Publishers. Pp.559-579
- Mitsui A, Enternmann B, Gill K (1983) Indoor and outdoor cultivation of *Tilapia* in seawater with algae as a sole food source. In Proceedings of the 2nd North Pacific Aquaculture System, Tokyo University, Japan, Pp. 323-340
- Anusuya. DM, Subbulaxmi G, Madhavi DM, Venkataraman LV (1981) 'Studies on the proteins of mass cultivated blue green alga (*Spirulina platensis*). J. Agric. Food Chem. 29:Pp522-525
- Gutiérrez-Rebolledo, G.A.; Galar-Martínez, M.; García-Rodríguez, R.V.; Chamorro-Cevallos, G.A.; Hernández-Reyes, A.G.; Martínez-Galero, E.(2015) Antioxidant effect of *Spirulina (Arthrospira) maxima* on chronic inflammation induced by Freund's complete adjuvant in rats. *J. Med. Food* , 18,Pp 865–871.
- Cohen Z (1986) 'Products from microalgae'. In *handbook of microalgal Mass Culture*. (Eds.). Richmond, A. Boca Raton FL, USA: CRC Press, Pp. 421-454
- Borowitzka MA (1994) In: Proceedings of the first Asia-Pacific Conference on Algal Biotechnology. (Eds.) Phang SM, Lee YK, Borowitzka MA and Whitton BA, University of Malaya, Kuala Lumpur, Pp. 5-15
- Glazer AN, Stryer L (1984) Phycofluor probes. *Trends Biochem. Sci.* 9:Pp423-427.
- Kronick MN (1986) The use of phycobiliproteins as fluorescent labels in immunoassay. *J Immunol Methods* 92:Pp1-13
- Rippka R, Deruelles J, Waterbury JB, Herdman M, Stainer RY (1979) Generic assignments. Strain histories and properties of pure cultures of cyanobacteria. *J. Gen. Microbiol.* 111: Pp1-61
- Fogg GE (1949) Growth and heterocyst production in *Anabaena cylindrica* Lemm. II. In relation to carbon and nitrogen metabolism. *Ann. Bot.* 13: Pp241-259
- Jacobson L (1951) Maintenance of iron supply in nutrient solutions by a single addition of ferric potassium ethylene diamine tetraacetate. *Plant Physiology* 26:Pp 411-413
- Allen MB, Arnon DI (1955) Studies on nitrogen fixing blue green algae. I. Growth and nitrogen fixation in *Anabaena cylindrica*. *Plant Physiol.* 30:Pp366-372
- Venkatarama, LV, Becker EW (1984) Production and utilization of the blue-green alga *Spirulina* in India. *Biomass* 4: Pp105-125
- Zarrouk C (1966) Contribution a l'étude d'une cyanophycee. Influence de divers facteurs physiques et chimiques sur la croissance et la photosyntheses de *Spirulina maxima*. Ph.D. Thesis. University of Paris, France
- Desikachary TV (1959) A monograph on Cyanophyta, ICAR Publication, New Delhi. Pp1-686
- Whyman M (1992) An *in vitro* method for estimation of phycoerythrin concentration in marine cyanobacteria (*Synechococcus* Sp.), *Limnol. Oceanogr* 37: Pp1300-1306
- Bennett A, Bogorad L (1973) complementary chromatic adaptation in a filamentous blue-green alga. *The Journal of Cell Biology* 58:Pp 419-435
- Olatz F, Ramon G, Juan LS (1991) Carotenoid composition in the Cyanobacterium *Phormidium laminosum*. Effect of nitrogen starvation. *FEBS Letters*, 282: Pp300-304
- Humberto J, Silva IC, Rodolfo JE (1989). Effect of nutritional factors on the culture of

Nostoc sp.as a source of phycobiliproteins.
Appl Microbiol Biotechnol 31:Pp 293-297
Vonshak A (1993) *Spirulina platensis*
(Arthrospira): Physiology, Cell Biology and
Biotechnology. Taylor &Francis, London.
ISBN 0-7484-0674-3
Roxana Olvera-Ramirez, Marisela Coria-Cedillo,
Rosa Olivia Canizares-Villanueva
Fernando Martinez-Jeronimo, Teresa Ponce-
Noyola, Elivira Rios-Leal,(2000). Growth
evaluation and byproduct characterization

of *Calothrix* sp. Bioresource Technology 72:
Pp121-124
Becker EW (1994) In: Microalgae -
Biotechnology and Microbiology, Cambridge,
Cambridge University Press, ISBN 0- 521
35020-4, Pp 1-4
Roe,S.(2000). Protein Purification Techniques:
A practical Approach.Oxford University
press, Oxford,UK.

Influence of different media on growth and phycobilins in *Phormidium fragile*

Sr.no	Medium	Fresh wt. (g)	Dry wt. (g)	Phycobilins %
1	BG-11	2.11±0.09 ^a	0.21±0.04 ^a	5.65±0.07 ^a
2	Allen & Amon	1.98±0.01 ^b	0.19±0.05 ^b	5.13±0.04 ^b
3	Fogg's Medium	1.70±0.00 ^c	0.12±0.01 ^c	5.61±0.06 ^a
4	Zarrouk' Medium	1.31±0.02 ^d	0.12±0.00 ^c	4.16±0.03 ^c
5	CFTRI	1.19±0.01 ^e	0.11±0.06 ^c	3.55±0.05 ^d