



## Influence of Sulphur Oxidizing Fungi on Growth and Yield of Soybean

**Kuldeepsingh Thakur, Kadali Satyakala, and Tini Pillai**

Plant Pathology Section, College of Agriculture Nagpur, Dr. Panjabrao  
Deshmukh Krishi Vidayapeeth, Maharashtra, India)

### Abstract:

Efficacy of two sulphur oxidizing fungal cultures viz., *Trichoderma harzianum* and *Aspergillus niger* individually and in combination with sulphur was evaluated in soybean crop in field trial at research farm of Plant Pathology Section, College of Agriculture, Nagpur, during *kharif*, 2012. The yield parameters viz., number of pods plant<sup>-1</sup>, grain yield and 1000 grains weight increased significantly due to inoculation of cultures under study with or without sulphur fertilizer. The treatment of *T. harzianum* and recommended dose of fertilizer (30:75:30 NPK kg ha<sup>-1</sup>) with sulphur (15 kg ha<sup>-1</sup>) had given maximum response in respect of number of pods plant<sup>-1</sup>, grain yield and 1000 grains weight. The number of pods plant<sup>-1</sup>, grain yield and 1000 grains weight were increased from 47.30 to 68.34, 823.00 to 1590.00 kg ha<sup>-1</sup> and 111.21 to 127.43 g by this treatment respectively over uninoculated control. The inoculation of *T. harzianum* + Sulphur (15 kg ha<sup>-1</sup>) + RDF could increase available sulphur content in soil from 13.36 kg ha<sup>-1</sup> in control to a maximum of 19.60 kg ha<sup>-1</sup> and sulphur uptake ranging from 250.00 to 322.33 mg 100g<sup>-1</sup> plant dry weight when compared with control.

### Key words:

Sulphur oxidizing fungi, yield, sulphur uptake, soybean.

### Introduction:

Soybean [*Glycine max* (L.) Merrill] is well known oilseed as well as pulse crop which is grown in various countries. Pulses and oilseeds are important constituents of Indian diet and supply a major part of the protein requirement. Soybean supplies 20 per cent oil and 40 per cent protein.

Sulphur is one of the seventeen essential nutrients and **it** is a structural component of amino acids, proteins, vitamins and enzymes and is essential to produce chlorophyll. Sulfur ranked equal to nitrogen for optimizing crop yield and quality. It enhances the efficiency of nitrogen for protein manufacture (Tucker, 1999). The major reservoir of sulphur in soil is





an unavailable elemental and reduced form of sulphur. Plants generally utilize the oxidised state of sulphate ( $\text{SO}_4^{2-}$ ) and therefore, it must first be oxidized before it can be used by crops. Most agricultural soils contain some micro organisms that are able to oxidize sulphur. The micro organisms are mainly responsible to make available sulphate from elemental or reduced forms of sulphur through its oxidation process with the intermediate formation of thiosulphate ( $\text{S}_2\text{O}_3^{2-}$ ) and tetrathionate ( $\text{S}_4\text{O}_6^{2-}$ ) in soils (Anandham and Sridar, 2004). Sulphur oxidizing fungi appear particularly useful for oxidation purpose since their spores or mycelia can be produced economically in large quantities and have good survival characteristics, both in the inoculants and in soil. Reports showed that combined inoculation of *Thiobacillus thiooxidans* and *Aspergillus niger*, *Trichoderma harzianum*, *Myrothecium cinctum*, *Aspergillus terreus* with sulphur improved soybean grain yield and protein content (Shinde *et al.*, 2004). Therefore, the study was undertaken to determine performance of sulphur oxidizing fungi on yield parameters of soybean, available sulphur in soil and on sulphur uptake.

### **Material and methods:**

The sulphur oxidizing fungi were isolated from soil samples collected from selected spots of different oilseed crops i.e. sesame, mustard, linseed and soybean fields of Agronomy Section, College of Agriculture, Nagpur. These isolates were further purified and screened for sulphur oxidizing ability in potato dextrose agar medium and czapek dox medium respectively. From these, two isolates i.e. *Aspergillus niger* and *Trichoderma harzianum* were selected for further experiment. A field experiment was conducted in randomized block design with three replications during *kharif*, 2012. There were nine treatments comprising of two fungal cultures and two sulphur levels (Table 1). The recommended dose of NPK (30:75:30 kg ha<sup>-1</sup>) was applied in the form of DAP and MOP, sulphur in the form of elemental sulphur powder. The cultures were multiplied in modified sulphur medium (Wainwright, 1978) and when they attained 10<sup>-7</sup> cfu ml<sup>-1</sup> strength, each one was mixed in 1:2 (v/w) carrier. Then each of the cultures @ 250 g 10 kg<sup>-1</sup> of





JS-335 seed of soybean were treated. The yield parameters of soybean viz., number of pods plant<sup>-1</sup>, grain yield and 1000 grains weight were studied. Five plants from each net plot were uprooted at harvest and all the developed pods were plucked and counted to estimate number of pods plant<sup>-1</sup>. The available sulphur content in soil and sulphur uptake by plant was analyzed by turbidimetric method (Chopra and Kanwar, 1980).

### **Result and discussion:**

The influence of sulphur oxidizing fungal cultures with and without sulphur on the growth and yield of soybean plant was found to be significant. Particularly, inoculation of *Trichoderma harzianum* culture along with sulphur (15 kg ha<sup>-1</sup>) showed significantly superior performance over all other treatments under study.

It is revealed from the data that number of pods plant<sup>-1</sup> were increased due to the cultures under study with and without sulphur over control. The maximum number of pods (68.34 plant<sup>-1</sup>) were recorded in the treatment *Trichoderma harzianum* culture and recommended dose of fertilizer along with sulphur (15 kg ha<sup>-1</sup>) and it was significantly superior over rest of the treatments and control. Next to this treatment application of *Aspergillus niger* culture and recommended dose of fertilizer along with sulphur (15 kg ha<sup>-1</sup>) significantly better than the remaining treatments except *Trichoderma harzianum* + recommended dose of fertilizer + Sulphur (10 kg ha<sup>-1</sup>). The remaining less effective treatments were viz., *Aspergillus niger* + recommended dose of fertilizer + Sulphur (10 kg ha<sup>-1</sup>), Sulphur (20 kg ha<sup>-1</sup>) + recommended dose of fertilizer, *Trichoderma harzianum* + recommended dose of fertilizer, *Aspergillus niger* + recommended dose of fertilizer, recommended dose of fertilizer in a descending manner. The minimum number of pods (47.30 plant<sup>-1</sup>) was recorded in uninoculated control. Similar trend of results was noticed in respect of grain yield. The maximum grain yield (1590 kg ha<sup>-1</sup>) was obtained in *Trichoderma harzianum* culture and recommended dose of fertilizer along with sulphur (15 kg ha<sup>-1</sup>) treatment and minimum weight of 1000 grains per treatment was also noticed to be the maximum (127.43 g) with the treatment of *Trichoderma*





*harzianum* culture and recommended dose of fertilizer along with sulphur ( $15 \text{ kg ha}^{-1}$ ). Shinde *et al.* (2004) in soybean reported that the number of pods, grain yield and 1000 grains weight were significantly increased due to combined inoculation of *Thiobacillus thiooxidans* and *Aspergillus niger*, *Trichoderma harzianum*, *Myrothecium cinctum*, *Aspergillus terreus* with sulphur. These results were conformed during the present studies so far as *T. harzianum* and *A. niger* are concerned. Shinde *et al.* (2000) in cotton reported that the number of picked bolls and seed yield were significantly increased due to combined inoculation of *Scolecobasidium constrictum*, *Myrothecium cinctum*, *Aspergillus terreus*, *Thiobacillus thioparus* and *T. thiooxidans*.

The available sulphur content in soil increased due to inoculation of sulphur oxidizing fungi in conjunction with sulphur application. The maximum available sulphur ( $19.60 \text{ kg ha}^{-1}$ ) was observed in the treatment of *Trichoderma harzianum* culture and recommended dose of fertilizer along with sulphur ( $15 \text{ kg ha}^{-1}$ ) and it was at par with *Aspergillus niger* culture and recommended dose of fertilizer along with sulphur ( $15 \text{ kg ha}^{-1}$ ) treatment. Next to these treatments the treatments were application of *Trichoderma harzianum* + recommended dose of fertilizer + Sulphur ( $10 \text{ kg ha}^{-1}$ ) and *Aspergillus niger* + recommended dose of fertilizer + Sulphur ( $10 \text{ kg ha}^{-1}$ ). The sulphur treatment alone also helped in increasing available sulphur level in soil from  $13.36 \text{ kg ha}^{-1}$  to  $17.40 \text{ kg ha}^{-1}$ . The less effective treatments were viz., *Trichoderma harzianum* + recommended dose of fertilizer and *Aspergillus niger* + recommended dose of fertilizer. The minimum available sulphur ( $13.36 \text{ kg ha}^{-1}$ ) was recorded in uninoculated control. Kadam *et al.* (2004) and Shinde *et al.* (2010) in soybean found increased available sulphur content in soil due to combined inoculation of *Thiobacillus thiooxidans* and *Aspergillus niger*, *Trichoderma harzianum*, *Myrothecium cinctum*, *Aspergillus terreus* with sulphur.

The data on sulphur uptake by soybean crop revealed that two fungal cultures with or without sulphur treatment increased sulphur uptake ranging from 250.00 to 322.33 mg  $100^{-1}$  g dry weight of plant and maximum was recorded in *Trichoderma harzianum* culture and recommended dose of





fertilizer along with sulphur (15 kg ha<sup>-1</sup>) and it was significantly superior over rest of the treatments. Next to this treatment the treatments were application of *Aspergillus niger* culture and recommended dose of fertilizer along with sulphur (15 kg ha<sup>-1</sup>), *Trichoderma harzianum* + recommended dose of fertilizer + Sulphur (10 kg ha<sup>-1</sup>) and *Aspergillus niger* + recommended dose of fertilizer + Sulphur (10 kg ha<sup>-1</sup>). The sulphur uptake also increased due to sulphur application alone from 250.00 to 262.00 mg 100<sup>-1</sup> g dry weight of plant. The less effective treatments were viz., *Trichoderma harzianum* + recommended dose of fertilizer and *Aspergillus niger* + recommended dose of fertilizer. The minimum sulphur uptake (250.00 mg 100<sup>-1</sup> g) was recorded in uninoculated control. Shinde *et al.* (2004) reported increase in sulphur uptake in soybean due to combined inoculation of *Thiobacillus thiooxidans* and *Aspergillus niger*, *Trichoderma harzianum*, *Myrothecium cinctum*, *Aspergillus terreus* with sulphur Earlier, Shinde *et al.* (2000) recorded similar results in groundnut and cotton crops.

**Table 1:** Effect of sulphur oxidizing fungi on yield parameters, available sulphur content in soil and sulphur uptake after harvesting of soybean crop.

S. No	Treatments	No. of pods plant <sup>-1</sup>	1000 grains Available		Sulphur	
			(kg ha <sup>-1</sup> )	weight (g)	sulphur content in soil (kg ha <sup>-1</sup> )	uptake (mg 100g <sup>-1</sup> )
T <sub>1</sub>	<i>Aspergillus niger</i> + RDF	53.00	1105.00	118.33	15.86	255.33
T <sub>2</sub>	<i>Trichoderma harzianum</i> + RDF	54.80	1180.00	119.51	16.06	258.66
T <sub>3</sub>	<i>A. niger</i> + Sulphur (10 kg ha <sup>-1</sup> ) + RDF	57.08	1272.33	121.21	17.89	283.66
T <sub>4</sub>	<i>T. harzianum</i> + Sulphur (10 kg ha <sup>-1</sup> ) + RDF	59.81	1374.66	123.01	18.42	289.33
T <sub>5</sub>	<i>A. niger</i> + Sulphur (15 kg ha <sup>-1</sup> ) + RDF	62.03	1440.33	126.67	19.00	318.00
T <sub>6</sub>	<i>T. harzianum</i> + Sulphur (15 kg ha <sup>-1</sup> ) + RDF	68.34	1590.00	127.43	19.60	22.33
T <sub>7</sub>	RDF (30:75:30 NPK kg ha <sup>-1</sup> )	55.33	1216.00	120.61	17.40	262.00
T <sub>8</sub>	Control	51.97	1045.00	115.16	15.06	53.33
T <sub>9</sub>		47.30	823.00	111.21	13.36	50.00
		1.51	32.54	0.70	1.14	3.39
		4.53	97.57	2.09	3.44	10.18





	SE $\pm$ (m) CD (P=0.05)		
--	-----------------------------	--	--

(Initial available sulphur content in soil is 13.03 kg ha<sup>-1</sup>)

### Conclusion:

The significant effect of different cultures on the performance of soybean may be attributed to the increased rate of microbial oxidation of elemental sulphur to sulphate (SO<sub>4</sub>) sulphur in soil which in turn might have increased the sulphur uptake and thereby the yield.

### References:

**Anandham, R. and R. Sridar, 2004.** Use of sulfur bacteria for increased yield and oil content of groundnut. In: Biofertilizers Technology, Kannaiyan, S., K. Kumar and K. Govindarajan (Eds.). Jodhpur, Scientific Publishers, India. pp. 365-371.

**Chopra, S. L. and J. S. Kanwar, 1980.** Analytical Agricultural Chemistry. Kalyani Publ., New Delhi. pp. 259-260.

**Kadam, R. M., D. B. Shinde, and A. C. Jadhav, 2004.** Effects of sulphur oxidizing microorganisms on nodulation pattern of soybean. J. Maharashtra agric. Univ. **29** (3): 300-302.

**Shinde, D. B., R. M. Kadam and A. C. Jadhav, 2004.** Effects of sulphur oxidizing micro-organisms on growth of soybean. J. Maharashtra agric. Univ. **29** (3):305-307.

**Shinde, D. B., A. C. Jadhav, R. M. Kadam, 2010.** Effect of sulphur oxidizing microorganisms on nutrient availability in soil under soybean. J. Maharashtra agric. Univ. **35** (1): 113-116.

**Tucker, M. R. 1999.** Essential plant nutrients: Their presence in North Carolina soils and role in plant nutrition. Agronomic Division, NCDA and CS: pp. 20-25.

**Wainwright, M. 1978.** A modified sulphur medium for the isolation of sulphur oxidizing fungi. Plant and Soil. **49**: 191-193.

