



HEAVY METAL CONCENTRATIONS IN WATER AND FISH *LABEO ROHITA* OF RIVER MULA, AT MULA DAM IN AHMEDNAGAR, MAHARASHTRA

M. J. SHAIKH

Department of Zoology, Arts, Commerce & Science College,
Sonai, Maharashtra, India
mishaikh1850@gmail.com

ABSTRACT:

The present study was conducted to determine the concentrations of heavy metals (Cd, Cr, Pb, Ni and Zn) in sample water and organs (muscle, gills and liver) of fish *Labeo rohita* of river Mula, at Mula Dam in Ahmednagar District were analyzed. There was an appreciable decrease in metal concentrations in sample water from site I to site II. The heavy metal concentrations in sample water was in the order Ni>Pb>Cd>Zn>Cr. Accumulation of heavy metals in the organs of fish *Labeo rohita* was found in the order gills>muscle>liver. In the gills, lead was found maximum (2.37 µg/g dry weight), whereas chromium was minimum (0.92µg/g dry weight). In the muscles, the values of lead was 1.98µg/g, comparatively chromium was 0.41µg/g. In the liver, lead, cadmium and nickel were more concentrated compared to zinc and chromium. The concentration of heavy metals accumulated in gills and liver was found in the order Pb>Cd>Ni>Zn>Cr and in the muscles, it was found in the order Pb>Cd>Ni>Cr>Zn. As the fish *Labeo rohita* is largely consumed by people, it is essential to make awareness about water pollution and effects of heavy metals on human body.

Keywords: Heavy metals, *Labeo rohita*, fish organs, Mula River, Mula dam.

INTRODUCTION:

The pollution of aquatic environment by inorganic and organic chemicals is a major factor posing serious threat to the survival of aquatic organisms including fish. Among the environmental pollutants, metals are of particular concern due to their potential toxic effect and ability to bio accumulate in aquatic ecosystems. The presence of heavy metals in aquatic ecosystem is the result of two main sources of contamination; natural processes and anthropogenic activities. Heavy metals including both essential and non essential elements have a particular significance in ecotoxicology, since they are highly persistent and all have the potential to be toxic to living organisms (Adeniyi and Yusuf, 2007).

Fish is often at the top of aquatic food chain and may concentrate large amounts of some metals from the water. Metal bioaccumulation is largely attributed to differences in uptake and depuration period of various metals in different fish species. Multiple factors including season, physical and chemical properties of water can play a significant role in metal accumulation in different fish tissues. The gills are directly in contact with water. Therefore, the concentration of metals in gills reflects their concentration in water where the fish lives, whereas the concentration in liver represents storage of metals in water (A.Rauf *et al.*, 2009). The studies carried out on various fishes have shown that heavy metals may alter the

physiological activities and biochemical parameters both in tissues and in blood (Vinodhini and Narayanan, 2008). Fishes are notorious for their ability to concentrate heavy metals in their muscles and since they play important role in human nutrition, they need to be carefully screened to ensure that unnecessary high level of some toxic trace metals are not being transferred to man through fish consumption.

The present research work aimed to analyze the accumulation of heavy metals (Cadmium, Chromium, Lead, Nickel and Zinc) in water and fish organs of *Labeo rohita*, of river Mula, at Mula Dam from Maharashtra.

MATERIALS & METHODS:

Water samples and fish samples (*Labeo rohita*) were collected from two different sites of river Mula in September 2015. Site I was near the wall of Mula Dam and Site II was 2 Km away from Dam, at upstream water of river Mula. Water samples were collected in the middle of the river at 50 cm below the surface, using 1 liter polythene bottles with screw caps. The bottle had been washed and soaked in 5% nitric acid and rinsed with de ionised water before use. The water samples were acidified immediately after collection by adding 5 ml nitric acid to minimize adsorption of heavy metals onto the walls of the bottles (APHA, 1998). Water samples were analyzed in Atomic Absorption Spectrophotometer for detection of heavy metals.

Samples of five fish (*Labeo rohita*) of nearly equal size and weight were dissected to remove muscles, gills and liver. The separated organs were put into petri dishes to dry at 120°C. The organs were placed into digestion flasks and ultrapure Con. Nitric acid and hydrogen peroxide (1:1 v/v) was added. The digestion flasks were then heated to 130°C until all the materials were dissolved. Digest was diluted with double distilled water appropriately. The heavy metals Cd, Cr, Pb, Ni and Zn were assayed using Atomic Absorption Spectrophotometer and the results were given as µg/g dry weight. Data obtained from the experiments were analyzed and the results were expressed as mean of all five. Values of $P < 0.05$ were considered statistically significant.

RESULTS & DISCUSSION:

Average heavy metal concentrations in the sample water collected from two sites of Mula Dam is given in Table – 1. The traces of heavy metals in water was found to decrease in the sequence Ni>Pb>Cd>Zn>Cr at both the sites. Maximum amount of metal found in sample water was nickel. The values of nickel observed were 5.84µg/l at site I and 5.26µg/l at site II. The values of chromium were minimum 1.03µg/l at site II and 1.27µg/l at site I. The higher level of lead, cadmium and zinc in the river water may be due to excessive use of pesticides and fertilizers in the agriculture fields along the banks of this river.

The analysis of heavy metals in different organs of fish *Labeo rohita* is shown in Table – 2. All the three organs – muscle, gills and liver shows different values of accumulation of heavy metals. Gills show higher level of metals succeeded by muscle and liver. In the gills, the sequence of trace metals is Pb>Cd>Ni>Zn>Cr. It is observed that the sequence of trace metals is Pb>Cd>Ni>Cr>Zn in muscles and

Pb>Cd>Ni>Zn>Cr in liver. The concentration of zinc and chromium shows fluctuations in all the three organs. The maximum amount of metals found in fish organs is lead, accumulating the gills. It shows values 2.37µg/g dry weights at site I and 2.12µg/g dry weights at site II. The minimum amount of heavy metal is chromium found in liver. It shows values of 0.26 µg/g in fishes collected from site I and 0.23 µg/g at site II. The amount of cadmium in gills is near to the amount of lead. The muscles show minimum amount of zinc (0.42 µg/g) and maximum amount of lead is 1.98 µg/g.

The values and sequence of heavy metals found in different organs are correlated with the results found by Abida Begum *et al.*, 2009. Accumulation of bioactive metals like cadmium, chromium, lead, nickel and zinc was actively controlled by fish through different metabolic processes and the level of accumulations usually depend on ambient concentrations. In the literature, heavy metal concentration in the tissues of freshwater fish vary considerably among different studies (Chattopadhyay *et al.*, 2002; Papagiannis *et al.*, 2004; V. S. Shrivastava *et al.*, 2011), possibly due to chemical characteristics of water, ecological needs, metabolism and feeding patterns of fish.

Table. 1- Average heavy metal concentrations (µg/l) in sample water collected from two sites of Mula Dam

Heavy metals	Site I	Site II
Cd	3.78	3.17
Cr	1.27	1.03
Pb	4.91	4.34
Ni	5.84	5.26
Zn	3.12	2.58

Table. 2- Average heavy metal concentrations (µg/g dry weight) in different organs of fish *Labeo rohita* collected from Mula Dam

Organ	Site I					Site II				
	Cd	Cr	Pb	Ni	Zn	Cd	Cr	Pb	Ni	Zn
Muscle	1.43	0.51	1.98	1.37	0.42	1.63	0.41	1.74	1.26	0.31
Gills	2.18	1.07	2.37	1.92	1.13	2.04	0.92	2.12	1.74	1.04
Liver	0.95	0.26	1.38	0.78	0.32	0.86	0.23	1.06	0.62	0.26

CONCLUSIONS:

The results of this study supply valuable information on the metal contents in water sample and fish organs of the river Mula, at

Mula Dam. Fish gills and muscles shows highest tendency to accumulate both lead and cadmium, while accumulation of metals is less in liver. As the fish *Labeo rohita* is one of the main aquatic

food, it should be noticed by the consumers and awareness must be done.

ACKNOWLEDGEMENT:

The author acknowledges sincere and heartfelt thanks to Principal, Arts, Commerce and Science College, Sonai for providing necessary laboratory facilities.

REFERENCES:

- Abida Begum, Hari Krishna S. and Irfanulla Khan, 2009.** Analysis of heavy metals in water, sediments and fish samples of Madivala lakes of Bangalore, Karnataka. *Int. J. Chem Tech Res.*, 1(2) : 245-249.
- Adeniyi, A.A. and K.A. Yusuf, 2007.** Determination of heavy metals in fish tissues, water and bottom sediments from Epe and Badagry Lagoons, Lagos, Nigeria. *Environ. Monitor. Assess.*, 37 : 451-458.
- APHA (American Public Health Association), 1998.** Standard methods for examination of water and waste water. 20th Edition, New York, USA.
- Rauf, M.Javed and M.Ubaidullah, 2009.** Heavy metal levels in three major carps (*Catla catla*, *Labeo rohita* and *Cirrhinamrigala*) from the river Ravi, Pakistan. *Pakistan Vet. J.*, 29(1): 24-26.
- Chattopadhyay, B., A. Chatterjee and S.K. Mukhopadhyay, 2002.** Bioaccumulation of metals in East Calcutta wetland ecosystem. *Aquat. Ecosys. Health Manag.*, 5 (2) : 191-203.
- Papagiannis, I., I. Kagalou, J. Leonardos, D. Petridis and V. Kalfakaou, 2004.** Copper and zinc in four freshwater fish species from Lake Pamvotis (Greece). *Environ. Int.*, 30 : 357-362.
- Vinodhini, R. and Narayanan, M., 2008.** Bioaccumulation of heavy metals in organs of fresh water fish *Cyprinus carpio* (Common carp). *Int. J. Environ. Sci. Tech.*, 5 (2) : 179 – 182.
- V. S. Shrivastava, Indrajit Sen and Ajay Shandil, 2011.** Study of determination of heavy metals in fish species of the River Yamuna (Delhi) by Inductively Coupled Plasma – Optical Emission Spectroscopy (ICP – OES). *Adv. Appl. Sci. Res.*, 2(2) : 161 – 166.