

Heavy Metal Pollution from a Thermal Power Station

R. D. Raut

J.B. College of Science, Ward ravishankar.raut@gmail.com

Abstract :-

The thermal power station situated near Chandrapur, India, releases fly ash into the environment and discharges effluent contaminated with fly ash and bottom ash through a main drain into the Erairiver. These fly ashes contain silicates, heavy metals and organic compounds. Thus, their discharge in the river affects the riverine water quality. In view of this the present study was undertaken. The concentration of heavy metal ions was determined in upstream and downstream of the river, thermal power station main drain and ash bund overflow to assess the heavy metal contamination level of the waters. Samples were collected from the selected sites at bi-monthly intervals over a period of one year and were analyzed for heavy metals Cu, Pb, Zn, Ni, Co, Cd, Mn and Cr using atomic absorption spectrophotometer. The study revealed that concentration of heavy metal ions was more in Erairiver downstream of the Thermal power station effluent drain, than upstream. It further revealed that the heavy metal content of the river water was within the standard permissible limits set for inland surface water.

Keywords :- heavy metal ions, effluents, ash bund, thermal power station, fly ash

Introduction:-

The use coal for the generation of electrical energy in the thermal power station produced large amount of ash which was generally disposed by using water to the nearby areas. According to recently compiled statistics, approximately 280 X 10⁶t of fly ash is Produce in the world every year by the combustion of coal¹ Coal fly ash consists primarily amorphous alumina silicates, heavy metals and organic compounds. Organic and inorganic constituents of coal ashes and slags may pose potential hazards to aquatic biota and humans². The concentration of heavy metals increases in water around the thermal power station due to release of fly ash into environment and discharge of effluents contaminated with fly ash and bottom ash³. Water quality around the thermal power station depends on the ash disposal methods, amount of suspended solids and heavy metals⁴. The purpose of this paper was to determine the quantity of heavy metals in effluents, in freshwater region of upstream of the Erairiver and number of spots downstream of the power station and also in the overflow of ash bund of thermal power station situated near Chandrapur, India.

Materials and Methods :-

Thermal power station is situated about 2 km towards north of Chandrapur. The gradation of coal used is E & F types. The Coal has calorific value of 3500-4500 kcal kg⁻¹. Its ash content is 30 to 40%, sulphur content 0.4 to 0.6% and moisture content 5 to 15%. Both units of the plant have modern electrostatic precipitator for precipitation of fly ash. The requirement of coal is nearly 7000t d⁻¹ and production of ash is nearly 2800t d⁻¹. This ash is transported in the form of slurry to ash bund. Out of this ash 5 t d⁻¹ in the form of fly ash is escapes from the electrostatic precipitor through chimney and spreads around an area of 1.32 km radius, depending on the wind velocity and direction.





Water samples were collected from the 4 selected sites at regular intervals over a period of one year, 5 time from each sites namely, (1) upstream and (2) downstream of Erai river, (3) power station main drain and (4) overflow of ash bund. The water samples from Erairiver were collected from 500m upstream and 500m down steam of discharge site of power station main drain of thermal power station. The samples were collected in 11 polythene bottles and were acidified with concentrate nitric acid and brought to the laboratory.

Each samples was then concentrated to 10ml volume by slow evaporation. The 20 ml of HNO_3 : H_2SO_4 (1:3) was added to each sample. It was evaporated to near dryness. The residue was extracted with 50ml double distilled water. Elements have been analyzed as per the standard methods recommended by APHA (1985). The concentration of heavy metals, Cu, Pb, Zn, Ni, Co, Cd, Mn and Cr were determined using flame atomic absorption spectrophotometer (GBC mode 906 AAS) by atomizing aqueous samples in air acetylene flame, Hollow cathode lamps for individual metals were used to produce respective resonance line.

Result and Discussion:

It was seen that concentration of heavy metals were within Bureau of Indian standards of permissible limits set for industrial effluents and drinking water (Table 1) The concentration of heavy metals. Cu, Pb, Zn, Ni, Co, Mn and Cr were in the following order ; power station main drian> ash bund overflow >Erai river downstream >river water upstream. In upstream water Co, Cd, and Cr were be law detectable limits and rest of the heavy metals were in very low concentration. The heavy metal level increased at downstream of Erai river. It is attributed to the disposal of effluents to the river by the thermal power station (Table. 1).

The power station main drain showed high variation in the concentrations of Pb, Ni, Co, Cd and Mn the ranges were 48- 180, 14-120, 94-156, 9-28 and 16-102, ug 1-1, respectively. Such higher concentration of heavy metals in power station main drain as compared to the ash bund overflow water was mainly due to the discharge of waster from water treatment plant, machine washing cooling tower blow down and fly ash. These values are lower than that reported for chandrapur Thermal Power Station (CPCB, 1988- 89).

In Erairiver downstream Ni and Cr were in very low concentration in the range of 6.0- 9.0 and 1.0-2.0 ug 1-1 respectively. The concentrations of Pb and Cd were high, ranging from 10.0- 25.30 and 3.0-4.0 ug 1-1, respectively, which exceeded maximum permissible limits prescribed by WHO (1996) : Pb =10 ug 1-1and Cd= 3ug 1-1, the rest of the heavy metals below recommended standard for drinking water . In all the samples except Kanhan river upstream, the concentration of Cd exceeded maximum permissible limit prescribed by WHO (1996) :Cd=3.0 ug 1-1 Higher concentration of Cd recorded in power station main drain was in the range of 9.0-28.0 ug 1-1, Which might be due to the discharge of effluents of water treatment plant and fly ash. The concentration of Pb in Erai river downstream, power station main drain and ash bund overflow was in the range of 10.0-25.0, 48.0-180.0 and 41.0 -52.0 ug1-1, respectively. In all the samples concentration of Pb exceeded maximum permissible limit 10.01-1, prescribed by





WHO (1996) . In the study area, concentration of Mn was found below prescribed maximum permissible limit, i.e. 500ug 1-1 (WHO,1996). In ash bund overflow the heavy metals were having low concentration as compared to power station main drain , it was due to percolation of ash water and setting of ash at the bottom of the ash bund (ISS, 1983).

Table- 1- Concentration of heavy metals in fresh and waster- water of Thermal power Station (ug1 -1) ND-Not detectable

Elements	Erai ri	iver	Erai	river	Power	station	Ash	bund
	upstream		downstream		main drain		overflow	
Copper	8.5-9.5		11.0 -15.0		2050.0		13-55.0	
Lead	5.7-6.4		10.0-25	5.0	48.0-18	30.0	41.0-52.	0
Zinc	3.0-4.4		11.0 -24	ŀ.0	12.0 -4	4.0	46.0-28.	0
Nickel	2.8 -3.6		6.0-9.0		14.0 -1	20.0	11.0 -15	.0
Cobalt	ND		5.0 -8.0		94.0-15	56.0	11.0 -17	.0
Cadmium	ND		3.0 -4.0		9.0-28.	0	3.0 -8.0	
Manganese	3.6 -4.6		9.0 -43.	0	16.0 -1	20.0	54.0 -92	.0
Chromium 🦾	ND		1.0-2.0	-	7.0-9.0)	6.0-8.0	

Conclusion:

The concentrations of all the heavy metal ions found in the area of power station main were within the bureau of Indian standards of permissible limits for industrial effluents prescribed by I.S.S. (1983). The concentration of heavy metals in the area of Erai river downstream was lower than the permissible limits prescribed by Indian standard (1983) but slightly higher as per WHO. This was attributed to the dilution effect of water. The water of Erai river is suitable for drinking purpose as per Indian Standard (1983) but not suitable as per WHO (1996) as its heavy metal content is slightly higher than its recommended standards. However, the water of Erai River is suitable for inland surface water and for irrigation purposes.

References:

1. Hockely, D. and Sloot H.A.V, Long term process in a stabilized coal waste block exposed to sea water. J. Envniron. Sci.Tech, 25: 1409. 1991

2. **Harris W.R.andSilberman, D.**thim dependent leaching of coal fly ash by cheletaating agents. J .Envniron. Sci.Tech, 17:139, 1983.

3. Nihan, C.T. Graydon J.W. and. Kirk D. W, Utilizing coal fly ash as a land fill barrier material. J. Waster Management, 16: 587-595. 1996

4. **Sarangi, P. K and Mishra A.P. C,** Soil activites and yield in ground nut, ladies finger and redish in fly ash amended soil . J. Res. Ur. Chem. Environ, 2 (2) :7-9. 1998

5. **Hegade, B.A. and Patil T.M.**Utilisation of industrial wastewater in agriculture. J. poll Res. 2:37- 39 1983.

6. Jain ,C.K.Bhatia K.S and Vijay T, Ground water quality in a coastal region of Andhra Pradesh. J. Indian Environ. Hith. 39: 182-192. 1997





7. Nair, J.C. and Ganapathi S, Water quality of the Bhadar river basin. J. Indian Environ. Hlth, 39: 197- 206. 1997

8. **APHA**, Standard methods for the examination of wastewater, Sixteenth edn, American Public Health association ,New York. 1985.

9. **Bureau Indian Standards,** Tolerance limites for industrial effluents. IS: 2490, Part I. 1981.

10. **CPCB**. Central Board for the Prevention and Control of Water Pollution, 1978 – 70 A assessment and development study of river basin series. Part- 1 Adsorbs- 13178- 70.

	Sr	Туре	of Membership	Subscription rate	
	1	Individ	ual life member	5000/-	
	2	Institutional life membership		10000/-	
r RTGS/ NEFT/ We Bank Name	stern N	Money Tr	ansfer/ Cash Deposit	our Bank Details are - NDIA	
Bank Name			STATE BANK OF I	NDIA	
ank Account Name			Vishwashanti Multip	urpose Society. Nagpur	
ccount No.			33330664869		
Account Type			Current		
FSC Code			SBIN0016098		
			SBININBB239		
Swift Code					
			16098		
Swift Code Branch Code MICR Code			440002054		

