



**CORYNANDRA CHELIDONII VAR. PALLAI (REDDY AND RAJU) V.S. RAJU; - NEW ADDITION  
 TO THE ANGIOSPERMIC FLORA OF MAHARASHTRA, INDIA.**

**SUBHASH R. SOMKUWAR<sup>1</sup>, JAGANNATH V. GADPAYALE<sup>2\*</sup> and ALKA CHATURVEDI<sup>3</sup>**

1. Department of Botany, Dr. Ambedkar College, Deekshabhoomi Nagpur-440 010.

2. Department of Botany, S. N. Mor College of Arts, Commerce and Smt. G. D. Saraf Science College, Tumsar (M.S.), India-441 912.

3. P.G.T. Department of Botany, University Campus, RTM University, Nagpur-440 033.

Corresponding author's E-mail: [gadpayale\\_jv@yahoo.co.in](mailto:gadpayale_jv@yahoo.co.in)

**ABSTRACT:**

Field survey resulted in collection of *Corynandra chelidonii* var. *pallai* (Reddy and Raju) V.S. Raju, of family Cleomaceae as a new addition to the flora of Maharashtra State with recent literature of classification system. The paper also envisages brief citation, description, flowering and fruiting time followed by ecological notes.

**Keywords:** *Corynandra chelidonii*, diversification, taxonomic literature and Phenology.

**INTRODUCTION:**

In the recent years there are some new additions were reported to the Flora of Maharashtra State such as *Juncus bufonius* L. (Lekhak et al., 2011), *Premna mollissima* Roth. (Paithane and Bhuktar 2013); *Hedychium flavescens* Carey ex Roscore (Jadhao et al., 2014); *Physalis pruinosa* L. (Somkuwar et al. 2014); *Amomum maximum* Roxb. And *Zingiber capitatum* Roxb. var. *elatum* (Roxb.) Baker. (Govekar and Sardesai 2015); *Opilia amentacea* Roxb. (Somkuwar et al. 2015); *Amorphophallus longiconnectivus* Bogner. And *Phyllodium longipes* (Craib) Schindl. (Gadpayale et al., 2016) etc.

The genus *Cleome* was first described by Linnaeus in *Species Plantarum*, Vol. 2 (1753: 671) of family Cleomaceae is almost universally accepted from the past 140 years (e.g. Hooker and Thomson 1872, Cooke 1903, Sundararaghavan 1993, Reddy and Raju 2001 and Kers 2003) with approximately 250 accepted species (Mabberley 2008 and The Plant List 2018) in the world (Cooke 1903).

Recently available literature of Iltis and Cochrane (2007, 2014) represents the division of *Cleome* s.l. into a number of segregate genera. According to Cochrane and Iltis (2014), the genus *Corynandra* Schrad.ex Spreng., based on *Corynandra chelidonii* (L. f.) Cochrane and Iltis (*Cleome chelidonii* L. f.)

which is re-established with other four combinations as *C. felina* (L. f.) Cochrane and Iltis, *C. flava* (Banks ex DC.) Cochrane and Iltis, *C. viscosa* (L.) Cochrane and Iltis and *C. viscosa* subsp. *nagarjunakondensis* (Sundararagh.) Cochrane.

In relation to this some worker as Russell L. et. al. (2017) in resolving generic boundaries in Indian-Australasian Cleomaceae: Circumscription of *Areocleome*, *Arivela*, and *Corynandra* as distinct genera and Feodorova, T. A. et al. (2010) provides molecular data for the diversification and the origin of C4 in *Cleome* (Cleomaceae) which also explore and concrete the opinions of Cochrane and Iltis (2014) which is represented by 18 species in India (Sundararaghavan 1993) and 12 species in Maharashtra state (Flora of Maharashtra state by BSI, Vol-I, 2000) including recent discovery of *Corynandra elegans* Chandore, U.S. Yadav and S.R. Yadav (2016).

In the present work *Cleome chelidonii* L. f. is now considered as *Corynandra chelidonii* (L. f.) Cocharane and Iltis. (2014) which is also consider by Subhash Sirangi and Ajmeera Ragan (2017) in their study of Dormancy and in-vitro seed germination of *Cleome chelidonii* var. *pallai* C.S. Reddy and V. S. Raju (2001) as *Corynandra chelidonii* var. *pallai* (reddy

and raju) V.S. Raju, ined. [(V.S. Raju, per. comm.) (2017)].

There is no any information in any regional flora and taxonomic literature about the occurrence of this species in Maharashtra State; hence this article is about the first time reporting of *Corynandra chelidonii* var. pallai (Reddy and Raju) V.S. Raju for the Angiospermic flora of Maharashtra State, India.

#### **METHOD AND MATERIAL:**

Field explorations were made in Eastern region of Maharashtra State, India to collect specimens in different seasons (2015 to 2018) in order to record the actual and essential key characters of specimens in reproductive and vegetative phases. Herbarium specimens were prepared and all relevant data were recorded in a field book along with photographs.

Taxonomic treatment for *Corynandra chelidonii* var. pallai (reddy and raju) V.S. Raju (SubhashSirangi and Ajmeera Ragan 2017; Reddy, C.S. and Raju, V.S 2001, a new variety of *Cleome chelidonii* L. f. Cleomaceae, Journal of Economic and Taxonomic Botany 25: 217–218).

#### **RESULTS AND DISCUSSION**

Erect, annual herb, much branched up to 1.5 m high; internodes to 12 cm. Basal leaves 3–5 foliate; leaflets up to 4-7 x 0.3-0.7 cm. lanceolate, cuneate at base, acute or rounded at apex margin crenate to crenate-serrate. Petiole 4- 8 cm, upper leaves 1–3 foliate; leaflets linear, base cuneate or rounded, apex acute, serrate. Petiole 2-4 cm. Flowers showy, pinkish or rosy in lax terminal corymbose racemes. Bracts foliaceous 1.5 mm long. Flower buds ovoid. Sepals elliptic, acuminate 1.5-2 mm. Petals ovate, 1.4- 1.7cm, rosy (pink). Stamens numerous (120–160); filaments 1.5- 1.8 cm, purple tinged. Ovary sessile, 8-10 mm long, glabrous. Capsule linear-cylindrical, narrow at the base, beak 5-10 mm long. Valves striate, parallel nerved, glabrous. Seeds

many asymmetrical ovoid, comma shaped, compressed. 1.3–1.8 mm in diam. dark brown, central portion paler and smooth, cleft narrow, slightly coloured, about 0.5- 0.7 mm deep. Testa with scattered scales, bearing waxy coating, elaisome absent.

#### **CONCLUSION:**

Habitat: Margins of Lake and Stream.


Distribution: Kondha, Bhuyar, Shahapur, Kardha, and Khapari (Chichal) Dist.- Bhandara (Maharashtra) India.

Phenology: Flowering and fruiting: July–December.

#### **REFERENCE :**

- Chandore, A N., U. S. Yadav and S. R. Yadav. 2016. A new elegant species of *Corynandra* (Cleomaceae) from Konkan region of Maharashtra, India. *Phytotaxa* 260 (1): 89–94.
- Cochrane, T.S. and Iltis, H.H. (2014) Studies in the Cleomaceae VII: Five New Combinations in *Corynandra*, an earlier name for *Arivela*. *Novon* 23: 21–26.
- Cooke, T. (1903) The Flora of the presidency of Bombay, Vol. 1. Taylor and Francis. London.
- Feodorova A., E. Voznesenskaya, G. Edwards and E. Roalson .2010. Biogeographic patterns of diversification and the origins of C4 in *Cleome* (Cleomaceae). *Systematic Botany* 35: 811–826. BioOne.
- Gadpayale J., Somkuwar S. R and AlkaChaturvedi. 2016. Two new addition to the flora of Maharashtra State, India. *Int. J. of Life Sciences, Special Issue A6*.
- Govekar Ravikiran and Sardesai M. M. (2015). Addition of Cardamom and Ginger for the Flora of Maharashtra. *Nelumbo the bulletin of the botanical survey of India, Vol. 57, ISSN-0976-5069*.
- Hooker J.D. and Thomson T. 1872. *Capparideae* In: Hooker, J.D. (Ed.) *The Flora of British India*. Vol. 1. Reeve and Co., London, pp. 167–180.

- Iltis, H.H. and Cochrane T.S. 2014. Studies in the Cleomaceae VI: A new genus and sixteen new combinations for the flora Mesoamericana. *Novon*23: 51–58.
- Iltis H.H. and Cochrane T.S. 2007. Studies in the Cleomaceae V: A new genus and ten new combinations for the flora of North America. *Novon*17: 447–451.
- JadhaoAnand, Vijay Paithane and Bhuktar A. S. 2014. A Ginger Lily *HedychiumFlavescens* Carey Ex Roscoe (Zingiberaceae): New records for Flora of Maharashtra, India. *Journal of Global Biosciences*. Vol. 3(5), pp. 792-794.
- Kers L.E. 2003. Capparaceae. In: Kubitzki, K. and Bayer, C. (Eds.) *The Families and Genera of Vascular Plants*. Vol. 5. Springer-Verlag, Berlin, pp. 36–56.
- Lekhak M. M., Chandore A. N. and S. R. Yadav. 2011. *Juncusbufonius* L. -An Addition to the flora of Maharashtra, India. *J. Bombay Nat. Hist. Soc.* 108 (1):68 – 69.
- Linnaeus C. 1753. *Species Plantarum*2: 671.
- Mabberley D. J. 2008. *The Plant-Book: A Portable Dictionary of the Vascular Plants* (Third Edition). Cambridge University Press, Cambridge, UK.
- Paithane Vijay and Bhuktar A. S. 2013. *Premnamollissima* Roth.-new reports for flora of Maharashtra (India). *The Indian Forester*, ISSN No. 0019-4816.
- Reddy C.S. and Raju, V.S. 2001. A new variety of *Cleome chelidoniif.* (Cleomaceae). *Journal of Economic and Taxonomic Botany* 25: 217–218.
- Russell L. Barrett, Eric H. Roalson, Kym Ottewell, Margaret Byrne, Sanjay P. Govindwar, Shrirang R. Yadav, Asif S. Tamboli, and Avinash R. Gholave. 2017. Resolving Generic Boundaries in Indian—Australasian Cleomaceae: Circumscription of *Areocleome*, *Arivela*, and *Corynandra* as Distinct Genera. *Systematic Botany* Vol. 42 (4) : 694-708, <https://doi.org/10.1600/036364417X696401>.
- Singh N.P. and Karthikeyan S. 2000. Flora of Maharashtra State: Dicotyledones, Vol. I (Ranunculaceae to Rhizophoraceae) Botanical Survey of India.
- Somkuwar S. R., Moghe R. P., Kamble R. B. and AlkaChaturvedi. 2015. *Opiliaamentacea* Roxb. Opiliaceae: A new genus record to the flora of Maharashtra, India. *Bioscience Discovery*, Vol. 6, Number 1 - I (Special).
- Somkuwar S. R., R. B. Kamble and Alka Chaturvedi. 2014. *Physalispruinosa* L. (Solanaceae): A new record to the Flora of Maharashtra, India. *life sciences leaflets* Vol.57,(Dec)2014.
- SubhashSirangi and Ajmeera Ragan. 2017. Dormancy and *In Vitro* Seed Germination of *Corynandrachelidonii* var. *Pallae*(Cleomaceae) An Endemic and Ethnomedicinal Herb. *International Journal of Pharmacy and Biological Sciences* Vol.7 (2): 240-244.
- Sundararaghavan, R. 1993. Capparaceae. In: Sharma, B.D. and Balakrishnan, N.P. (Eds.) *Flora of India*. Vol. 2. Botanical Survey of India, Calcutta, pp. 248–335.
- The Plant List. 2018. Version 1.1. <http://www.theplantlist.org/tp1.1/search?q=cleome> (accessed 2 August 2018).



**FIG.1- *Corynandra chelidonii* var. *pallai* (Reddy & Raju) V.S. Raju; i-Root view, ii- in habitat, iii- Flowering, iv- single twig with buds, v- flowering mode at node, vi- single flower, vii- fruit.**