



EFFICACY OF SOLVENT EXTRACTS OF SOME MEDICINAL PLANTS FOR POTENTIAL ANTIBACTERIAL ACTIVITY.

Chandrashekhar J. Chandekar and Priya L. Lokhande

Shri Shivaji Science College, Congress Nagar, Nagpur-12

Corresponding author Email : chandekarc@yahoo.com

Abstract

Abstract Different solvent extracts of the leaves of *Acacia Arabica*, *Ailanthus excelsa*, *Alstonia scholaris*, *Callistemom lanceolantus*, *Casurina equisetifolia*, *Catharanthus roseus*, *Grevellia robusta*, *Moringa oleifera*, *Psidium guava*, *Saraca indica*, *Tamarindus indicum*, *Vitex negundo* and *Thuja compacta* were investigated for their antibacterial activity on *Escherchia coli* NCIM-2931, *Proteus vulgaris* NCIM-2857, *Pseudomonas auriginosa* NCIM-5029, *Staphylococcus aureus* NCIM-5021, *Bacillus cereus* NCIM-2155, *Bacillus subtilis* NCIM-2063, and *Bacillus megaterium* NCIM-2087. 50 µl of solvent extract is used to test antibacterial activity using agar well diffusion method with well size of 5 mm diameter using High sensitivity test agar medium. Acetone and Methanol extracts of *Acacia Arabica*, *Callistemom lanceolantus*, *Casurina equisetifolia*, *Psidium guava*, *Saraca indica*, and *Thuja compacta* showed significant activity against tested bacteria. *Psidium guava* extracts in acetone, chloroform, methanol and petroleum ether, are active against all bacteria tested except *Bacillus cereus*. *Saraca indica*'s extract in acetone, chloroform, methanol and petroleum ether are active against all Gram positive bacteria tested.

Keywords

Key words: Solvent extracts, Antibacterial activity, agar well diffusion method.

Introduction

Introduction Medicinal plants form the backbone of traditional system of medicine in India. Pharmacological studies have acknowledged the value of medicinal plants as potential source of bioactive compounds (Prusti et.al., 2008). Phytochemicals from medicinal plants serve as lead compounds in drug discovery and design (Chakravarthy and Gode, 1985; Ebi and Ofoefule, 2000). Medicinal plants are rich source of novel drugs that forms the ingredients in traditional system of medicines, modern medicines, nutraceuticals, food supplements, folk medicines, pharmaceutical intermediates, bioactive principles and lead compounds in synthetic





drugs(Ncube,2008). Nowadays multiple drug resistance has developed due to the indiscriminate use of commercial antimicrobial drugs commonly used in the treatment of infectious disease(Davis,1994; Service,1995).In addition to this problem, antibiotics are sometimes associated with adverse effects on the host including hypersensitivity, immune-suppression and allergic reactions(Ahmad et.al.,1998).This situation forced scientists to search for new antimicrobial substances.Given the alarming incidence of antibiotic resistance in bacteria of medical importance(Monroe and Polk,2000),there is a constant need for new and effective therapeutic agents(Bhavani and Ballow 2000).Therefore ,there is a need to develop alternative antimicrobial drugs for the treatment of infectious diseases from medicinal plants (Clark,1996 and Cordell,2000).Several screening studies have been carried out in different parts of the world.There are several reports on the antimicrobial activity of different herbal extracts in different regions of the world(Chung et.al.,2004,De Boer et.al.,2004,Nair and Chanda,2004,Nair et.al.,2005)

Material and Method

Materials and Methods Test Organisms Escherichia coli NCIM 2931, Proteus vulgaris NCIM 2857, Pseudomonas aureginosa NCIM 5029, Bacillus cereus NCIM 2155, Bacillus subtilis NCIM 2063 , Bacillus megaterium and staphylococcus aureus NCIM 5021 these cultures were obtained from the National Collection of Industrial microorganisms, Pune. Plants Acacia arabica , Ailanthus excelsa, Alstonia scholaris, Casurina equisitifolia, Catharanthus roseus, Callistemom lanceolantus, Grevellia robusta, Moringa oleifera ,Psidium guayava, Saraca indica, Tamarindus indicum, Thuja compacta and Vitex negundo. Aqueous extraction Ten grams of dried plant material was extracted with 100 ml of distilled water for 6 h.at slow heat.Every 2 h.it was filtered through 8 layers of muslin cloth and centrifuged at 5000 g for 15 min.The supernatant was collected.This procedure was repeated twice and after 6 h.the supernatant was concentrated to make the final volume one fifth of the original





volume(Parekh,2005).The extract was then autoclaved at 1210C and 15 lbs pressure,and stored at 40C. Solvent Extraction Ten grams of dried plant material was extracted with 100 ml of solvent kept on a rotary shaker for 24 h.Thereafter,it was filtered and centrifuged at 5000 g for 15 min.The supernatant was collected and the solvent was evaporated to make the final volume one-fifth of the original volume(Parekh,2005).It was stored at 40C in airtight bottles for further studies. Antibacterial assay The antibacterial activity of different plant species was evaluated by agar well diffusion (Perez et.al.,1990)for solvent extracts using Hi-sensitivity test agar(M 485) for the assay.The microorganism was activated by inoculating a loopful of the strain in 5 ml of Hi-sensitivity test broth and broth was incubated at 350C+ 0.50C for 18-20 hrs.After incubation a loopful of actively growing culture was inoculated into 10 ml of Hi-sensitivity broth. Broth was incubated at 350C+0.50C for 6-8 hours.0.5 ml inoculums was inoculated in to the Hi-sensitivity test agar media and after proper homogenization it was poured into 100 mm petri dishes. For the agar well diffusion, a well was made in the seeded plates with the help of a cup-borer (5 mm).The test compound was introduced in to the well and all the plates were incubated at 350C+ 0.50C for 24 hrs. Microbial growth was determined by measuring the diameter of the zone of inhibition are presented (Table-1).

Result and Discussion

Results and Discussions Results are given in Table No.1.Successful prediction of botanical compounds from plant material is largely dependent on the type of solvent used in the extraction procedure. Traditional healers use primarily water as the solvent but in our studies we found that plant extracts in organic solvent (Acetone, Chloroform, Methanol and Petroleum ether) provided more consistent antimicrobial activity compared to those extracted in water. These observations can be rationalized in terms of the polarity of the compounds being extracted by each solvent and, in addition to their intrinsic bioactivity, by





their ability to dissolve or diffuse in the media used in the assay. *Ailanthus excelsa*, *Alstonia scholaris*, *Grevillia robusta*, *Moringa oleifera* (only water extract is active), *Tamarindus indica* extracts were totally inactive against all the microorganisms tested. The plants that exhibited antibacterial activity to a certain degree were *Acacia Arabica*, *Callistemom lanceolantus*, *Casurina equisetifolia*, *Psidium guajava*, *Saraca indica* and *Thuja compacta*. Acetone and methanol extract of *Acacia Arabica* were active against all the bacteria except *E. coli*. Methanol extract of *Callistemom lanceolantus* is active against *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Bacillus cereus*, *Bacillus subtilis* and *Bacillus megaterium*. Acetone, methanol and petroleum ether extracts of *Psidium guajava* are active against all bacteria (Fig. 1 and 2) except *Bacillus cereus*. Acetone, chloroform, methanol and petroleum ether extracts of *Saraca indica* are active against *Staphylococcus aureus* (Fig. 4), *Bacillus cereus*, *Bacillus subtilis* (Fig. 3), and *Bacillus megaterium* whereas water extract is active against *Bacillus subtilis*, and *Bacillus megaterium*. Acetone extracts of *Thuja compacta* is active against all Gram positive bacteria. Chloroform extract is active against *Proteus vulgaris*, *Bacillus subtilis* (Fig. 5), and *Bacillus megaterium* (Fig. 6). Methanol and petroleum ether extracts are active against *Staphylococcus aureus*, *Bacillus subtilis*, and *Bacillus megaterium* whereas water extract is inactive against all the microorganisms. Acetone and chloroform extracts of *Vitex negundo* are active against *Bacillus subtilis*, and *Bacillus megaterium*. Methanol extract is active against *Proteus vulgaris*, *Bacillus subtilis*, and *Bacillus megaterium*. Petroleum ether and water extracts are inactive against all the microorganisms. Out of 13 plant species, *Callistemom lanceolantus*, *Psidium guajava*, *Saraca indica* and *Thuja compacta* showed significant activity. The plant extracts were more active against the Gram positive bacteria. This is in agreement with previous reports that plant extracts are more active against Gram positive than against Gram negative bacteria (Vlietinck et al., 1995, Rabe et al., 1997). From our investigation of screening different plant species, the results obtained confirm





the therapeutic potency of some plants used in traditional medicine. The results of the present study support the folkloric usage of the studied plants and suggest that some of the plant extracts possesses compounds with antibacterial properties that can be used as antimicrobial agents in new drugs for the therapy of infectious diseases caused by pathogens. The most active extracts can be subjected to isolation of the therapeutic antimicrobials and undergo further pharmacological evaluation. Screening of various natural organic compounds and identifying active agents is the need of the hour, because successful prediction of lead molecule and drug like properties at the onset of drug discovery will pay off later in drug development.

Conclusion

From our investigation of screening different plant species, the results obtained confirm the therapeutic potency of some plants used in traditional medicine. The results of the present study support the folkloric usage of the studied plants and suggest that some of the plant extracts possesses compounds with antibacterial properties that can be used as antimicrobial agents in new drugs for the therapy of infectious diseases caused by pathogens. The most active extracts can be subjected to isolation of the therapeutic antimicrobials and undergo further pharmacological evaluation. Screening of various natural organic compounds and identifying active agents is the need of the hour, because successful prediction of lead molecule and drug like properties at the onset of drug discovery will pay off later in drug development.

Acknowledgement

Authors are indebted to Dr. D.K.Burghate, principal sses science college, congress nagar, Nagpur for their kind support during the research work. Thankfulness is also due to Dr. A.D.Bobdey, Prof. R.N. Deshmukh and Prof V.A.Dakhole for their constant encouragement during my research work.





Reference

- Ahmad I, Mehmood Z, Mohammad F, 1998. Screening of Some Indian medicinal plants for their antimicrobial properties. 1998. *J Ethnopharmacol* 62:183-193.
- Bhavani SM, Ballow CH, 2000. New agents for Gram positive bacteria. *Curr Opin Microbiol* 3:528-534.
- Chakravarty BK and Gode KD, 1985. Isolation of epicatechin from *Pterocarpus marsupium* and its pharmacological action. *Planta Medica* 1:56-59.
- Chung PY, Chung LY, Ngeow YF, 2004. Antimicrobial activities of Malaysian plant species. *Pharm Biol* 42:292-300.
- Clark AM, 1996. Natural products as resource for new drugs. *Pharm Res* 13:1133-1141.
- Cordell GA, 2000. Biodiversity and drug discovery a symbiotic relationship. *Phytochemistry* 55:463-480.
- Davis J. Inactivation of the antibiotics and the dissemination of resistance genes, 1994. *Science* 264:375-385.
- De Boer HJ, Kool A, Broberg A . 2005. Antifungal and antibacterial activity of some herbal remedies from Tanzania. *J Ethnopharmacol* 96:461-469.
- Ebi GC and Ofoefule SI, 2000. Antimicrobial activity of *Pterocarpus osun* stems. *Fitoterapia* 71:433-435.
- Nair R, Chanda SV, 2004. Antimicrobial Activity of Some medicinal plants of Saurashtra region. *J Tissue Res* 4:117-120.
- Nair R, Kalariya T, Chanda S, 2005. Antimicrobial Activity of Some selected Indian medicinal flora. *Turk J Biol* 29:41-47.
- Parekh J, Nair R, Chanda S, 2005. Preliminary screening of some folkloric plants from western India for potential antimicrobial activity. *Indian J Pharmacol*.





- Perez C,Paul M,Bazerque P.1990.Antibiotic assay by agar-well diffusion method.Acta Biol Med Exp 15:113-115.
- Prusti A,Mishra SR,Sahoo S and Mishra SK,2008.Antimicrobial Activity of Some Indian Plants. Ethnobotanical Leaflets 12:227-230.
- Rabe T, Van Staden J.1997.Antibacterial activity of South African plants used for medicinal purposes.J Ethnopharmacol 56:81-87.
- Vlietinck AJ,van Hoof L,Totte J et.al.1995.Screening of hundred Rwandese medicinal plants for antimicrobial and antiviral properties.J Ethnopharmacol 46:31-47.

