



BIOAUGMENTATION PACKAGE OF EGG SHELL BASED BIOSURFACTANT

FORMULATION : AN INNOVATIVE APPROACH FOR MONOCROTOPHOS

DEGRADATION IN SOIL

Kirti V. Dubey

Sevadal Mahila Mahavidyalaya, Nagpur

Corresponding author Email : kirtivijay_dubey@yahoo.com

Abstract:

The ability of the biosurfactant producing *Kocuria turfanesis* to mineralize monocrotophos was investigated through an experimental set up using different treatments of soil with carrier based bioaugmentation package of egg shell coated with biosurfactant and biosurfactant producing microbial isolates BS-J cells. Dual capabilities of biosurfactant production and degradation of monocrotophos are the two main attributes of strain BS-J which was isolated from lube oil and distillery spent wash contaminated soil collected from a distillery unit. On the basis of the cellular morphology, physiological and chemotaxonomic characteristics and phylogenetic similarity of 16S rDNA gene sequences, the strain BS-J was identified as a *Kocuria turfanesis*. Present study has shown that degradation of monocrotophos in contaminated soil was facilitated after using carrier based bioaugmentation package of egg shell coated with biosurfactant producing microbial isolate BS-J cells and its biosurfactant resulted 86-89 % degradation of monocrotophos after 7-days of treatment. Without biosurfactant producing microbial cells the degradation of pesticide was comparatively lower indicating the need of live biosurfactant producing microbial cells and their surfactants for efficient degradation of pesticide. Results indicate that *Kocuria turfanesis* strain BS-J has great potential utility for the bioremediation of soil contaminated with monocrotophos.

Keywords: Biosurfactant, biodegradation, curd whey, monocrotophos, soil.

Introduction:

Monocrotophos was amongst the top 15 pesticides used in the 20th century. Monocrotophos works systemically and, on contact, it is highly toxic to organisms, including humans (Isoda et al., 2005). Monocrotophos [di-methyl-(E)-1,2-methylcarbamoylevinylphosphate] is an organo-phosphorus insecticide. It has been widely used to control a variety of insects on crops, such as cotton, sugar cane, peanuts and tobacco, because of its low cost and effectiveness.





(Sha, 1999). Monocrotophos is weakly sorbed by soil particles because of its hydrophilic nature. Leaching of monocrotophos may pollute the surface and/or groundwater, ultimately resulting in adverse effects on biological systems (Subhas & Singh, 2003). In general, pesticide degradation in soil can be influenced by both biotic and abiotic factors, which act in tandem and complement one another in the microenvironment (Singh et al., 2003). Microbial activity has been deemed to be the most influential and significant cause of organo-phosphorus pesticide removal. Therefore, biodegradation is a reliable and cost-effective technique for pesticide abatement, and a major factor determining the fate of organo-phosphorus pesticides in the environment (Kertesz et al., 1994; Munnecke & Hsieh, 1974). Monocrotophos is characterized by a P–O–C linkage and amide bond, and has been reported to be degraded as a sole carbon or phosphorus source in liquid media by *Pseudomonas aeruginosa* sp., *Clavibacter michiganense* ssp. (Subhas & Singh, 2003), *Arthrobacter atrocyaneus* sp., *Bacillus megaterium* sp. and *Pseudomonas mendocina* (Bhadbhade et al., 2002a, b, c). The majority of studies concerning the fate of monocrotophos in soils have focused on tropical soil systems (Racke et al., 1996; Vijay et al., 2006). There have been no reports of the degradation of monocrotophos in tropical soils in presence of microbial surfactants. Hence, a study was undertaken to determine the ability of *Kocuria turfanesis* strain BS-J to produce biosurfactant in curd whey and to utilize the whole fermented curd whey that contains the cells and biosurfactant for degradation of monocrotophos in contaminated soils. In the present work, studies were conducted to determine the degradation of monocrotophos in different treatments of soil with carrier based bioaugmentation package of egg shell coated with biosurfactant and biosurfactant producing microbial isolates BS-J cells. Carrier based bioaugmentation process using biosurfactant was chosen with an aim to provide sustainable degradation of pesticide in soil.





Material and Methods:

Screening of biosurfactant producing microorganisms Biosurfactant producing microorganisms were isolated from soil collected from an area just below the spent wash pumping device of a distillery unit which was contaminated with lube oil and distillery spent wash by using selective enrichment procedure and plating serially diluted enriched culture on sterile nutrient agar followed by incubation at 37°C for isolated colonies (Dubey and Juwarkar, 2001). Isolates so obtained were individually screened for biosurfactant production from processed curd whey waste on the basis of stability of foam, emulsification index, surface tension reduction, and biosurfactant yield as per the methods described by Dubey and Juwarkar, 2001. Among five isolates tested, BS-J was an efficient isolate with potential of having high emulsification index indicating powerful biosurfactant which can be used to solubilize / mobilize pesticide in contaminated ecosystem and produce high yield of biosurfactant in curd whey. Identification of the efficient biosurfactant producing isolate Biosurfactant producing microbial isolate designated as BS-J was identified on the basis of morphological, cultural and biochemical methods described by Cappuccino and Sherman, 1999 and the results were compared with those in Bergey's Manual of Systemic Bacteriology (Holtz et al., 1994). 16S rRNA gene amplification and sequencing of biosurfactant producing isolate BS-J The 16S rRNA gene amplification and sequencing of biosurfactant producing microbial isolate BS-J was performed by the following procedure: DNA was extracted by using ZR fungal/bacterial genomic DNA extraction kit (Zymo Research Corporation 17062 Murphy Ave. Irvine, CA 92614, U.S.A.) according to the manufactures instructions and used for PCR amplification of 16S rRNA gene. The 16S rRNA gene was amplified by PCR using universal bacterial primers 6-27f (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492 r (5'-TACGGYTACCTTGTTACGACTT-3'). The amplified product was electrophoresed on 1% agarose gel and a band corresponding to 1.5 Kb was cut and was eluted using QIA quick PCR purification Kit (Qiagen, Hilden, Germany). This purified





amplicon was then sequenced using Big Dye Terminator cycle sequencing kit and an ABI PRISM model 3130xl analyzer automatic DNA sequencer (Applied Biosystems, USA) with 3 different internal sequencing primers (27f, 535r and 1492r). The sequences obtained were aligned, edited manually and the aligned sequence was used for BLAST search. Alignment of 16S rRNA sequence and construction of phylogenetic tree of biosurfactant producing isolates The sequence of 16S rRNA gene was aligned with closely related sequences using CLUSTAL_X Windows interface (Thompson et al., 1997) and edited manually. Neighbour-joining analysis was done with Kimura 2-parameter model (Kimura, 1980), using TREECON; version 1.3b (Van de Peer and De Watcher, 1997). The stability among the groupings of phylogenetic tree was assessed by taking 1000 replicates. For J-strain *Kytococcus sedentarius* (X87755) was used as an out-group. Strain J formed a clade with *Kocuria turfanesis* Ho-9042T. Based on nucleotide homology and phylogenetic analysis, the biosurfactant producing BS-J was identified as *Kocuria turfanesis* at IMTECH, Chandigarh, India, and MTCC allotted number was 10635. Enrichment of biosurfactant producing *Kocuria turfanesis* strain BS-J for monocrotophos degradation: *Kocuria turfanesis* strain BS-J is a biosurfactant producing microbial culture isolated from lube oil and distillery spent wash contaminated soil collected from a distillery unit and was added to 500 ml Erlenmeyer flasks containing 100 ml of synthetic medium which consisted of (g/l) NaNO₃, 2.0; K₂HPO₄, 1.0; KH₂PO₄, 0.5; MgSO₄.7H₂O, 0.5; KCl, 0.1 and FeSO₄.7H₂O, 0.01 amended with 50 ppm (wv-1) Monocrotophos as sole carbon source. The microbial cultures were subjected to selective enrichment with sequential and weekly transfer of strain BS-J to an increasing concentrations of Monocrotophos from 50 - 500ppm. Inoculum preparation for degradation of monocrotophos in mineral salts medium studies: Strain BS-J was pre-cultured in baffled Erlenmeyer flasks containing LB medium. Flasks were incubated overnight at 30 oC on a rotary shaker at 150 rpm. The contents of the inoculated flasks were centrifuged at 8000 rpm for 10 min and the cell pellet was washed three times with fresh





medium and quantified by the dilution plate count technique. For all experiments, 10^6 CFU ml⁻¹ was used and samples were incubated at 30°C at 150 rpm unless otherwise stated. Determination of monocrotophos in cell free fermented mineral salts medium After periodic intervals of 24 hours the fermented broth was centrifuged at 8000 rpm for 10 min and the supernatant samples were collected and filtered through a 0.45 micron membrane filter paper. Amount of monocrotophos in the filtered samples was estimated by method of Janghel et al. 2006. Monocrotophos degradation kinetics of *Kocuria turfanesis* For the study of degradation kinetics of *Kocuria turfanesis* strain BS-J pure culture of MCP1 was suspended in 1 ml 0.9% saline to make a cell suspension of 1×10^8 cells per ml and 100 µl of this suspension was inoculated in 125 ml of M1 medium containing 150 mg L⁻¹ concentration of monocrotophos and incubated at 28 ± 2 °C for 15 days in an orbital shaking incubator at 90 rpm under aerated culture conditions. Thereafter, monocrotophos was extracted at time interval 5, 10 and 15 days twice with equal amount of ethyl acetate (1:1). The solvent was evaporated and the residue was re-dissolved in 3 ml of ethyl acetate. Amount of monocrotophos was estimated at 254nm. The residual amount of monocrotophos was calculated by molar absorbion coefficient. Degradation kinetics of monocrotophos follows first order kinetics. Degradation constant K_{deg} was calculated by drawing straight line curve between log of concentration of monocrotophos at certain incubation duration Vs incubation duration and the slope of the curve was multiplied by 2.303. Thus, the degradation coefficient was calculated by using straight line equation:- $y = -mx + c$ $K_{deg} = -2.303 \times -m$ Half life of monocrotophos in M1 medium by MCP1 was calculated by the following formula- $t_{1/2} = 0.699/K_{deg}$. The value was expressed in hours. Degradation of monocrotophos in soil and its estimation Experimental setup for monocrotophos degradation in soil: Garden soil with sand, mixed in 2:1 proportion were filled in experimental polypropylene pots at the rate of 1 Kg in each pot and following treatments were given to the soil to study the role of





biosurfactant produced by Strain BS-J in monocrotophos degradation periodically after 5 days of incubation in green house at a temperature of 30 oC over a period of 30 days. 1. Soil amended with monocrotophos + 10% tap water 2. Soil amended with monocrotophos + 10% curd whey 3. Soil amended with monocrotophos at the rate of 500 ppm + 10% fermented curd whey containing biosurfactant +whole cells of BS-J 4. Soil amended with monocrotophos at the rate of 500ppm + 10% cell free fermented curd whey containing biosurfactant During incubation distilled water was added to adjust the moisture content to 40% of the maximum water-holding capacity. The samples were incubated at 30 oC in the dark. Subsamples were extracted periodically extracted at time interval of 5 days over a period of 30 days, twice with equal amount of ethyl acetate (1:1). The solvent was evaporated and the residue was re-dissolved in 3 ml of ethyl acetate. Amount of monocrotophos was estimated at 560 nm using UV-Vis spectrophometer by method of Janghel et al. 2006.

Result and Discussion:

Collection of curd whey wastes for biosurfactant production and their physico-chemical characteristics Industrial waste viz. curd whey is a viable alternative source for biosurfactant production and it has been already reported (Dubey and Juwarkar, 2001). However, in the present study was carried out with an aim to replace mineral salts medium with curd whey so that no-cost medium for biosurfactant production by strain BS-J can be assessed for its use in remediation of pesticide contaminated soil. Results presented in Table 1 show that curd whey i.e. lactic acid whey had high COD of 56000 mg/l , sugar and nitrogen levels of 6.8 g/l and 987.0 mg/l respectively indicating that it had sufficient organic load required for the growth of biosurfactant producing microbial isolate. Curd whey is generated during the preparation of Chakka which is an intermediate product obtained by draining of curd (a type of fermented milk from lactic acid bacteria) for preparation of “Shrikhand”, a well known popular Indian dessert. During preparation of 1 Kg Chakka, 9 l o lactic





acid whey is generated (Bandyopadhyay and Mathur, 1987). Suitability of curd whey as a fermentation medium for biosurfactant production was assessed by studying the growth profile of biosurfactant producing isolate and its biosurfactant production potential. Use of curd whey in the present study has been taken up instead of using sweet whey as it is feasible to transport the curd whey to the biosurfactant production site without using any special cryo-preservation techniques because continued fermentation of curd whey by indigenous lactic acid bacteria of curd will lead to more production of lactic acid which is a preferred substrate for biosurfactant production since the biosurfactant producer used in this study is an efficient utiliser of lactic acid for growth (Holtz et al., 1994). Isolation and screening of efficient biosurfactant producing microorganism The use of biosurfactants is an attractive option because of its versatility, biodegradability, ecological safety and environmental acceptance. However, their high production cost limits their use in bioremediation processes. In this context, it is necessary to evaluate the culturing conditions that optimize their production, assess the economic use of new substrates, such as those arising from industrial waste, and to evaluate techniques of isolation and purification to make production more economically feasible. In order to form a basis for using no-cost wastes as potential alternative fermentative medium for biosurfactant production, present study focused on the use of industrial waste viz. curd whey as nutrient medium for biosurfactant production by new microbial isolate. For isolation and screening of biosurfactant producing microorganisms, the phenomenon of reduction of surface tension of the culture medium and emulsification index was selected as described earlier (Dubey and Juwarkar, 2001). The success of biosurfactant production depends on the development of cheaper processes and the use of low-cost raw materials, which account for 10-30% of the overall cost (Rodrigues et al., 2006; Makkar et al., 2002). A great variety of agro-industrial wastes have been studied as potential substrates for biosurfactant production such as by-product of the sugar cane industry, fruit processing industry, whey wastes





represents an alternative medium for the biosurfactant production process as these have no-cost as compared to other known substrate sources, and they possess valuable nutrients required for the fermentation process (Dubey and Juwarkar, 2001 and 2004, Dubey et al., 2005). Results presented in Table 2 show that BS-J isolate had biosurfactant production capacity which was evident by production of stable foam formation in the fermented curd whey waste that lasted up to two hours of standing. The isolate reduced the surface tension of the fermented curd whey waste from an initial range of 56 mN/m to 27 mN/m indicating the production of effective biosurfactant and also showed good emulsification property as the emulsification index E24 of 94% was obtained. Biosurfactant yield produced by isolate was 0.98 g/l. Results have shown that the isolate produced high yields of biomass and biosurfactant in curd whey owing to the presence of very rich source of organic carbon, nitrogen and minerals like calcium, phosphorus, potassium, sodium, copper and iron. It is also a good source of vitamins of B-complex group viz. riboflavin and pantothenic acid that are readily available for the growth of biosurfactant producing isolates (Nickerson, 1974 Reductions in COD and total nitrogen, sugars, and phosphate contents were observed in the whey waste which indicates decrease in pollutional load of the waste during biosurfactant production. Identification of efficient biosurfactant producing strain: Based on morphological, biochemical, physiological characteristics, the efficient biosurfactant producing isolate BS-J was identified as *Kocuria turfanensis*. This biosurfactant producing isolate is a new strain as revealed by 16S rRNA sequence pattern and phylogenetic tree analysis performed at IMTECH, Chandigarh, India, and MTCC allotted number to this isolates as 10635 (Dubey et al. 2011). Utilization of monocrotophos and biosurfactant production by isolate BS-J in mineral salts medium Monocrotophos degradation by isolate BS-J was monitored by for a period of 120 h. After 24 h, 50% of pure monocrotophos had rapidly disappeared, followed by a slower decrease of monocrotophos with longer incubation times. During 72 to 96 h of incubation





the surface tension of the cell free broth dropped from 62 mN/cm to 28 mN/cm indicating the production of biosurfactant which was maximum at 96 h of incubation with an yield of 0.58g/l. The degradation of monocrotophos supported cell growth, indicating that isolate BS-J could utilize monocrotophos as a carbon source. Such an isolate with dual potential of producing biosurfactant along with monocrotophos degradation can be used an advanced approach for the bioremediation of wastewater or soil contaminated with monocrotophos. From the culture enrichment study it was evident that the concentration of monocrotophos affects the growth of BS-J. The optimal concentration of monocrotophos for the growth of BS-J was 200 mg l⁻¹, and a concentration higher than 600 mg l⁻¹ was toxic for the normal growth of BS-J isolate. Growth kinetics of *Kocuria turfanesis* under optimized culture conditions In order to ascertain the viability of *Kocuria turfanesis* strain BS-J in monocrotophos containing mineral medium, the growth kinetics was studied. Application of 200 mg L⁻¹ concentration of monocrotophos at pH-7, temperature 28 °C to the strain BS-J resulted in visible increase in the turbidity in the medium. It was observed that under optimized culture conditions and in the presence of monocrotophos the growth of strain BS-J was 0.550mg ml⁻¹ and it was 0.243mg ml⁻¹ in monocrotophos free medium. There was an increase of 51 % in the growth indicating that monocrotophos induced the growth of strain BS-J by serving as a source of phosphorus and energy. Degradation kinetics of *Kocuria turfanesis* In order to find out the half life of monocrotophos in the presence of strain BS-J , the degradation kinetics was studied in mineral salts medium. Results have shown that monocrotophos degradation kinetics correlated well with its growth kinetics. It was found that more the growth of strain BS-J, higher was the rate of monocrotophos degradation. This could be due to utilization of monocrotophos by strain BS-J as sole source of phosphorus and additional energy source during its growth period. After 7 days of incubation the recovery of monocrotophos was 13% i.e 26.2 mg l⁻¹ from its initial applied concentration 200 mg l⁻¹ and the





percentage of degradation of monocrotophos was found to be approximately 90%. The degradation constant K_{deg} calculated from straight line curve was 0.00817 and half -life was 79.32 h. In control sample the kinetic constant K_{deg} was zero and thus the half life was infinite indicating that strain BS-J is an efficient degrader of monocrotophos. Degradation of monocrotophos in different treatments of soil with carrier based bioaugmentation package of egg shell coated with biosurfactant and biosurfactant producing microbial isolates BS-J cells The experimental set up used to study the degradation of monocrotophos in different treatments of soil with carrier based bioaugmentation package of egg shell coated with biosurfactant and biosurfactant producing microbial isolates BS-J cells is presented in Figure 1. Physico-chemical characteristics of soil used in the present study is presented in Table 3 . Results show that soil had pH of 6.9 with bulk density of 1.16 g/cm³ and water holding capacity of 61.04%. Organic carbon content of soil was 0.56%. The N, P and K levels of soil were 0.52, 0.073 and 0.144, respectively. There was decrease in the pH of the soil from 6.9 to 6.4 after treatment with 500 ppm monocrotophos due to acidic nature of the pesticide. However, after amendment with 10% egg shell coated with curd whey the pH of the monocrotophos treated soil improved to 7.8-7.9 which is due to calcium carbonated nature of the egg shell that was used as a carrier material for coating biosurfactant for facilitated degradation of monocrotophos in contaminated soil. This improvement in pH of soil is very important to maintain the physiological state of the native microbial population present in the contaminated soil that maintains the biogeochemical cycle in soil. This is owing to the fact that microorganisms are a major component of the ecosystem and play a considerable role in the degradation of insecticides. This study indicates that use of calcite nature of the egg shell can act as soil conditioning agent. It has been proven that addition of biosurfactants, bioemulsifiers, and/or biosurfactant-producing microorganisms can be used in soil biodegradation techniques, soil washing, and water and waste treatment (in situ and ex situ) (Urumand Pekdemir, 2004; Zhou and Zhu, 2008).





Biosurfactants have also been found to be useful for oil spill remediation and for dispersing oil slicks into fine droplets and converting mousse oil into an oil-in-water emulsion (Toledo et al., 2008). In the present study, degradation of monocrotophos in contaminated soil was found to be facilitated after using carrier based bioaugmentation package of egg shell coated with biosurfactant and biosurfactant producing microbial isolate BS-J cells and its biosurfactant resulted 86-89 % degradation of monocrotophos after 7-days of treatment. Without biosurfactant producing microbial cells the degradation of pesticide was comparatively lower indicating the need of live biosurfactant producing microbial cells and their surfactants for efficient degradation of pesticide. In general, most pesticides used in agriculture are moderately hydrophobic compounds, with complex molecular structures that differ from hydrocarbons in their lower hydrophobicity and in the presence of a polar functional group. These compounds are also strongly adsorbed by soil organic matter and desorption is limited (Rodríguez-Cruz et al., 2004). Wattanaphon et al. (2008) evaluated the ability of a BS biosurfactant produced by *Burkholderia cenocepacia* BSP3 to enhance pesticide solubilization for further application in environmental remediation. The results showed that the application of the BS biosurfactant to facilitate pesticide solubilization demonstrated that this biosurfactant at concentrations below and above its CMC could enhance the apparent water solubility of methyl parathion, ethyl parathion and trifluralin. In the present study the chemical composition of soil in terms of total organic carbon, nitrogen, phosphorus and potassium also improved with the addition of the bioaugmentation package which facilitates the biodegradation of pesticide. The efficiency of degradation in the active soil samples (without inoculation) was generally better than that in the sterile soil samples (without inoculation), suggesting that microorganisms in the soil may play a role in the degradation of monocrotophos. The best degradation of monocrotophos was in the soil samples with the addition of the BS-J, indicating that BS-J isolate from lube oil and distillery spent wash contaminated soil collected from a distillery





unit can compete and survive with the local microflora in the soils. The result of monocrotophos degradation in soils proved that BS-J could be used successfully for the removal of monocrotophos from contaminated soils. This is the first report on a Kocuria turfanesis strain BS-J which, showed the combined capabilities of producing biosurfactant from monocrotophos as the carbon source and also degradation of monocrotophos in soil. Earlier our studies have shown fairly stable surface active properties of biosurfactant produced by Kocuria turfanesis strain BS-J in regard to emulsification of pesticides viz. monocrotophos and imidacloprid used routinely for agricultural purposes at extremes of different environmental conditions (Dubey et al. 2012).

Table 1: Characteristics of curd whey collected for biosurfactant production

S. No.	Type of waste	Sources of collection of wastes	Parameters					
			pH	COD (mg/l)	BOD (mg/l)	Total sugars (g/l)	Total Nitrogen (mg/l)	Total Phosphorus (mg/l)
1.	Whey waste (WW)	Amruta Dairy, Umred	4.3	56,000	28,000	6.8	987.0	352.0

Findings are the mean values of the three replicate readings

Table 2: Variation in biosurfactant production potential and other related parameters of the new microbial isolate BS-J using curd whey as fermentation medium (After 120 hours of incubation)

Parameters	Control	Microbial isolate BS-J
Foaming	-ve	+ve
Emulsification index (%)	-	94
Surface tension (mN/m)	56	27
Ph	7.0	8.57
Biomass yield (c.f.u./ml)	12x10 ²	83x10 ⁸
COD (mg/L)	37000	19340
Biosurfactant yield (g/l)	0.0011	0.9897

Results are the mean values of the three replicate readings






Table 3. Physico-chemical characteristics of Monocrotophos contaminated soil and degradation of monocrotophos in different treatments of soil with carrier based bioaugmentation package of egg shell coated with biosurfactant and biosurfactant producing microbial isolates BS-J cells

Different treatments of the soil sample	Bulk density (g/cm ³)	Maximum water holding capacity (%)	pH	Organic carbon (%)	Nitrogen (%)	Phosphorus (%)	Potassium (%)	% degradation of monocrotophos
Soil + monocrotophos (500 ppm) + 10% tap water	1.16	60.20	6.6	0.63	0.059	0.083	0.156	7.12
Soil + monocrotophos (500 ppm) + 10% egg shell coated with curd whey	1.13	60.34	6.5	0.75	0.073	0.089	0.167	21.45
Soil + monocrotophos (500 ppm) + 10% egg shell coated with fermented curd whey containing biosurfactant + BS-J cells	1.12	60.48	7.8	0.79	0.08	0.092	0.180	88.72
Soil + monocrotophos (500 ppm) + 10% egg shell coated with cell free fermented curd whey containing biosurfactant produced by BS-J	1.14	60.45	7.9	0.77	0.07	0.090	0.178	85.71

Results are the mean values of the three replicate readings



Figure 1: Flow sheet of an experimental setup for monocrotophos degradation in soil in presence of biosurfactants adsorbed on egg shell as carrier material



1

2

Conclusion:

The carrier based technology used in the present study for the cleanup of pesticide contaminants in soil, included bioaugmentation package of egg shell coated with biosurfactant, and microbe. This unique formulation stimulates and enhances the bioremediation processes designed to disperse and augment remediation of pesticide in soil. Carrier based biosurfactant technology for pesticide decontamination is a simple, highly effective, simple to use, cost-effective and completely non-toxic, environmentally friendly, as all its ingredients are organic in origin and completely biodegradable. This bioaugmentation package of egg shell coated with biosurfactant and biosurfactant producing microbial isolate BS-J resulted 86-89 % degradation of



monocrotophos after 7-days of treatment. Application of egg shell waste as a carrier of biosurfactants that improves the physico-chemical and microbiological status of disturbed soil thereby improving the overall fertility and rejuvenation of disturbed soil ecosystem.

Acknowledgement:

Author acknowledges the financial support of University Grants Commission, New Delhi for carrying out this work under Project No. F. No. 38-199/2009(SR) for carrying out this work under Major Research Project. Author also acknowledges the support of Management and the Principal of the college to provide infrastructure required to carry out the present study.

References:

- Bandhoapdhy AK, Mathur B N (1987) Indian milk products: a compendium. Dairy India 211-218.
- Bhadbhade BJ, Dhakephalkar PK, Sarnaik SS & Kanekar PP (2002a) Plasmid-associated biodegradation of an organophosphorus pesticide, monocrotophos, by *Pseudomonas mendocina*. Biotech Lett 24: 647–650.
- Bhadbhade BJ, Sarnaik SS & Kanekar PP (2002b) Bioremediation of an industrial effluent containing monocrotophos. Curr Microbiol 45: 346–349.
- Bhadbhade BJ, Sarnaik SS & Kanekar PP (2002c) Biomineralization of an organophosphorus pesticide, Monocrotophos, by soil bacteria. J Appl Microbiol 93: 224–234.
- Cappuccino JG, Sherman N (1999) Microbiology: A laboratory manual. Addison-Wesley Longman, Inc., Harlow, England. pp. 199-204.





- Dubey, K. and Juwarkar, A. (2001). Distillery and curd whey wastes as viable alternative sources for biosurfactant production. *World J. Microbiol. Biotechnol.*, 17: 61-69.
- Dubey, K. V., Juwarkar, A. A., (2004). Determination of genetic basis for biosurfactant production in distillery and curd whey wastes utilizing *Pseudomonas aeruginosa* strain BS2. *Indian J. Biotechnol.* 3, 74-81.
- Dubey, K. V., Juwarkar A. A., Singh, S. K., (2005). Adsorption-desorption process using activated carbon for recovery of biosurfactant from distillery waste. *Biotechnol. Prog.* 21, 860-867.
- Dubey K. V., Charde, P. N. Meshram, S. U. Yadav, S. K. Singh S. K. and Juwarkar A. A. (2011). Biosurfactant Production Potential of New microbial isolates using Combination of Distillery Waste with Other Industrial Wastes. 'Special Issue of 'Microbial Biosurfactants /Marine Microbial Biosurfactants ' International Open Access Journal of Petroleum and invironmental Biotechnology (DOI:<http://dx.doi.org/10.4172/2157-7463.S1-002>).
- K. V. Dubey, P. N. Charde, L. P. Shendre, S. U. Meshram, V. S. Dubey and A. A. Juwarkar (2012) Surface-active Properties of Novel Biosurfactants at Extremes of Environmental Conditions useful in Remediation of Pesticides Contaminated Soils. *Bioresource Technology* 126:368-374.
- Holtz JG, Krieg NR, Sneath PHA, Staley JT & Williams ST (1994) *Bergey's Manual of Determinative Bacteriology*, 9th edn. Williams & Wilkins Co., Baltimore. Isoda H, Talorete TP, Han J, Oka S, Abe Y & Inamori Y (2005) Effects of organophosphorus pesticides used in China on various mammalian cells. *Environ Sci* 1: 9-19.
- Janghel E. K, Rai, . J. K. , Rai M. K and Gupta V. K. (2006) A new and highly Sensitive spectrophotometric determination of monocrotophos in





- environmental, agricultural and biological Samples . Journal of the Chinese Chemical Society 53:343-347.
- Kertesz MA, Cook AM & Leisinger T (1994) Microbial metabolism of sulfur- and phosphorus-containing xenobiotics. FEMS Microbiol Rev 15: 195–215.
- Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. Journal of Molecular Evolution 16:111-120.
- Makkar RS, Cameotra SS (2002) An update on the use of unconventional substrates for biosurfactant production and their new applications. Appl Microbiol Biotechnol 58:428–434.
- Munnecke DM & Hsieh DPH (1974) Microbial decontamination of parathion and p-nitrophenol in aqueous media. Appl Environ Microbiol 69: 5198–5206.
- Nickerson TK (1974) Lactose: Fundamentals of Dairy Chemistry. Webb, B. H. Johnson, A. M., Alford, J. A. (Edited) Westport, Connecticut A VI, pp. 273-324.
- Racke KD, Steele KP, Yoder WA, Dick WA & Avidov E (1996) Factors affecting the hydrolytic degradation of chlorpyrifos in soil. J Agric Food Chem 44: 1582–1592.
- Rodrigues LR, Teixeira JA, van der Mei HC, Oliveira R (2006) Physicochemical and functional characterization of a biosurfactant produced by *Lactococcus lactis* 53. Colloids and Surfaces B: Biointerfaces 49:79–86.
- Sha JJ (1999) A Manual of Industrial and Chemical Production: Agricultural Chemicals, pp. 13–14. Chemical Industry Press, Beijing. Singh BK, Walker A, Morgan JAW & Wright DJ (2003) Effects of soil pH on the biodegradation of chlorpyrifos and isolation of a chlorpyrifos-degrading bacterium. Appl Environ Microbiol 69: 5198–5206.





- Subhas & Singh DK (2003) Utilization of monocrotophos as phosphorus source by *Pseudomonas aeruginosa* F10B and *Clavibacter michiganense* subsp. *Insidiosum* SBL 11. *Can J Microbiol* 49: 101–109.
- Rodríguez-Cruz, M.S., Sánchez-Martín, M.J., Sánchez-Camazano, M. (2004). Enhanced desorption of herbicides sorbed on soils by addition of Triton X-100. *J. Environ. Qual.* 33, 920-929.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin, F, Higgins DG (1997) The CLUSTAL_X Windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 22: 4673-4680.
- Toledo, F.L., González-López, J., Calvo, C. (2008). Production of bioemulsifier by *Bacillus subtilis*, *Alcaligenes faecalis* and *Enterobacter* species in liquid culture. *Bioresour. Technol.* 99, 8470-8475.
- Van de Peer Y, De Wachter R (1997). Construction of evolutionary distance trees with TREECON for Windows: accounting for variation in nucleotide substitution rate among sites. *Computational and Applied Biosciences* 13: 227-230.
- Wattanaphon, H.T., Kerdsin, A., Thammacharoen, C., Sangvanich, P., Vangnai, A.S. (2008). A biosurfactant from *Burkholderia cenocepacia* BSP3 and its enhancement of pesticide solubilization. *J. Appl. Microbiol.* 105, 416-423.
- Vijay AKB, Gundi VA & Reddy BR (2006) Degradation of monocrotophos in soils. *Chem* 62: 396–403.
- Urum K, Pekdemir T.(2004) *Chemosphere*. Evaluation of biosurfactants for crude oil contaminated soil washing. *Dec*;57(9):1139-50.
- Zhou, W., Zhu, L. 2008. Enhanced soil flushing of phenantrene by anionic-nonionic mixed surfactant. *Water Res.* 42, 101-108.

