



## USE OF EXTRACTIVE SPECTROPHOTOMETRY AND ION-PAIR FORMATION WITH CONGO RED FOR THE DETERMINATION OF PROMETHAZINE HYDROCHLORIDE ANTIPSYCHOTIC DRUG IN PHARMACEUTICAL FORMULATION

D. M. CHAFLE<sup>1</sup>


<sup>1</sup>Department of Chemistry, Taywade College, Mahadula-Koradi, Nagpur, Maharashtra, India- 441 111  
[dmchafle@gmail.com](mailto:dmchafle@gmail.com)

### Abstract:

Promethazine hydrochloride is phenothiazine used as antipsychotic drug. A simple, rapid and sensitive extractive spectrophotometric method is described for the determination of promethazine hydrochloride in pure form and in its pharmaceutical formulation. The method is based on the ion-pair complex formation between the drug and the Congo red dye in acidic medium. The red coloured chromogen, being extractable with chloroform, could be measured quantitatively at 483 nm. All variables were studied to optimize the reaction conditions. The method obeys Beer's law and the precision and accuracy of the method were checked. The molar composition and stability constant of the ion-pair complex formed is determined at room temperature by Turner and Anderson method. The molar composition of complex Congo red : PromethazineHCl is 1:2. The stability constant is  $1.2489 \times 10^4$ . The detection limit is 19.4 to 83.5  $\mu\text{g ml}^{-1}$  and the molar absorptivity is  $3.5298 \times 10^3 \text{ L mol}^{-1} \text{ cm}^{-1}$  and Sandell's sensitivity is 0.1125  $\mu\text{g cm}^{-2}$  per 0.001 absorbance unit.

**Key words:** Extractive spectrophotometry, ion-pair complex, promethazine hydrochloride, congo red.

**INTRODUCTION:** Phenothiazines are widely used drugs in the treatment of mental disease. Promethazine hydrochloride is one of the phenothiazine which shows neuroleptic, antihistaminic and antibacterial activity. It is chemically 2-RS-N,N-dimethyl-1-(10Hphenothiazine-10 yl)propan -2- amine hydrochloride. The structure of promethazine hydrochloride is as shown below.



**Figure 1: Structure of Promethazine**

It is official in B.P. [1], USP [2] and I.P. [3]. Literature survey reveals that various methods for its determination are reported which include flow injection analysis [4, 5], chromatography [6, 7], spectrofluorometry [8, 9], solid phase extraction [10]. The spectrophotometric method [11, 12] involved the chromogenic reaction with various reagents. Few extractive spectrophotometric methods [13-15] has been reported for the estimation of promethazine hydrochloride. The objective of the present study was to develop a simple economic extractive spectrophotometric method with easily available chromogenic reagent Congo red which can be used for routine quality control of drug in bulk and in its formation.

### EXPERIMENTAL

**Instrument :** A Equiptronic model EQ282 visible spectrophotometer with a pair of 1 cm matched quartz cells was used for optical measurements.

### Materials :

**Standard stock solution :** A 0.01M stock solution of pure promethazine hydrochloride (PMTH)(Sigma) was prepared in 0.1 N hydrochloric acid solution. From this stock solution working standard solution of 32  $\mu\text{g/ml}$  was prepared by appropriate dilutions.

**Congo red (CR) solution:** A 0.01 M congo red solution was prepared in 0.1 N hydrochloric acid. The reagent solution of desired concentration was then obtained by further dilutions.

All chemicals used were of analytical grade. Chloroform and water used were double distilled.

### METHOD:

**Calibration curve:** In a series of flasks an appropriate aliquots of standard PMTH solution were added and final volume was made to 15 ml. These aliquots were vigorously shaken for 15 minutes in two installments of 5 ml chloroform. The absorbance of one of the extracted layer was first scan in visible range for determination of maximum wavelength,  $\lambda_{\text{max}}$  (Figure 2). At this wavelength then absorbance for all other solutions of extracted chloroform were measured against reagent blank and calibration curve was plotted for validation of Beer's Law (Figure 3).

**Stability constant:** 0.001M and 0.002M solutions of PMTH and CR were prepared by

appropriate dilution. Different aliquots of drug and chromogenic reagent were mixed in varying proportion of mole fraction of drug. These aliquots were then similarly extracted with chloroform as discussed above. A Turner and Anderson [16] plot was plotted (Figure 4) to calculate stability constant of the complex formed.

**Estimation of drug in tablet:** Commercial tablet Phenargan were procured from local market. Twenty tablets were weighed and ground to fine powder in mortar. An amount of powder equivalent to 10 mg of drug was transferred to 100 ml volumetric flask. 0.1 N HCl was added and shaken thoroughly to dissolve the powder completely. The volume was made up to the mark and solution was filtered. Suitable aliquots were mixed with CR solution and extracted with 10 ml chloroform in two installments of 5 ml. Absorbance of extracted chloroform was obtained at  $\lambda_{max}$ . The amount of drug in tablet was then calculated from calibration curve. The optical characteristic and results of error analysis is given in Table 1 and Table 2.

#### RESULT AND DISCUSSION

Congo red and promethazine hydrochloride forms ion pair complex in acidic medium. The drug and congo red individually are insoluble in water while the complex formed is insoluble in water and highly soluble in chloroform. This makes the basis of the proposed extractive spectrophotometric method.

The red coloured ion-pair complex formed shows broad absorption spectra with  $\lambda_{max}$  483 nm (Figure 2). In Turner and Anderson plot (Figure 4) the maxima obtained at mole fraction = 0.666 of drug. Hence the molar composition of the complex formed is found to be PMTH : CR = 2:1. The stability constant was also calculated from same plot (Figure 4) by using equation

$$K = \frac{x}{(a_1 - x)(b_1 - x)} = \frac{x}{(a_2 - x)(b_2 - x)} \text{ and } x = \frac{a_1 b_1 - a_2 b_2}{(a_1 + b_1) - (a_2 + b_2)}$$

It is found to be  $1.2489 \times 10^4$ . The Beer's Law plot is found to be satisfactorily linear over the range of  $19.4 \mu\text{g ml}^{-1}$  to  $83.5 \mu\text{g ml}^{-1}$ . The ion-pair complex is intensely red coloured shows molar absorptivity  $3.5298 \times 10^3 \text{ L mol}^{-1} \text{ cm}^{-1}$ . The recovery study of drug shows good percentage recovery of 99.794% with relative standard deviation of 0.1907 and standard deviation of  $\pm 0.01903$ .

#### CONCLUSION:

The proposed method is simple, accurate and reproducible and can be used for routine

quality control analysis of promethazine hydrochloride.

#### REFERENCES

- [1] British Pharmacopoeia,(1993): H.M. Stationary Office, London. Pp 522.
- [2] The United State Pharmacopoeia, (1970): 18<sup>th</sup> Revision, mac Publishing Co. Eastern PA. Pp 280.
- [3] Pharmacopoeia of India, (1985): Ministry of Health & Family Welfare, 3<sup>rd</sup> ed., Govt. of India. Pp 420.
- [4] Caltayud J.M., Mateo V.G. (1992): Cerium (IV) arsenite as a solid-phase reactor for use in flow-injection analysis. Spectrophotometric determination of promethazine. *Analytica Chimica Acta*. 264 (2) : Pp. 283-289.
- [5] Caltayud J.M., Sancho T.G. (1992): Spectrophotometric determination of promethazine by flow injection analysis and oxidation by Ce (IV). *Journal of Pharmaceutical and Biomedical Analysis*. 10 (1) : Pp. 37-42.
- [6] Sreenivasa R.V., Sivrama P. K., Lakshmi P., Bourne D. W. A., (2001): A liquid chromatographic method for the simultaneous determination of promethazine and three of its metabolite in plasma using electrochemical and UV detectors. *Journal of Chromatographic Science*. 39 (2) : Pp. 70-72.
- [7] Laitem L., Bello I., Gaspar P. (1978): Gas chromatographic determination of tranquilizer residues in body fluids and in the meat of slaughtered animals. *Journal of Chromatography A*. 156 (2): Pp. 327-329.
- [8] Homyak I., Kozma L., Lapat A., Tovari i. (1997): Spectrofluorimetric determination of diethazine in pharmaceutical preparations. *Biomedical Chromatography*. 11 (2): Pp. 99-101.
- [9] Szydłowska- Czerniak A., Dembinski B., Kurzawa M. (2001): Spectrofluorometric determination of 2- and 10-disubstituted phenothiazines. *Acta.Pol. Pharm*. 58 (4): Pp. 235-240.
- [10] Song Q. , Putcha L. (2001): Quantization of promethazine and metabolites in urine samples using on-line solid-phase extraction and column-switching. *J. Chromatogr B Biomed Sci. Appl*. 763 (1-2): Pp. 9-20.
- [11] Gowda B.G., Seethramppa J., Melwanki M.B., (2003): Spectrophotometric determination of antiallergic drug in bulk powder and in its pharmaceutical formulations. *Journal of Analytical Chemistry* 58(6): Pp. 509-513.
- [12] Devani M.B., Suhagia B. N., Shah S. A. (1999): Spectrophotometric determination of

promethazine hydrochloride in bulk and in its dosage forms. Indian Journal of Pharmaceutical Sciences. 61(2): Pp. 110-112.

[13] Basavaiah K., Krishnamurthy G. (1998): Extractive spectrophotometric determination of some phenothiazine derivatives in pharmaceutical preparations. Talanta. 46 (4): Pp. 665-670.


[14] Rewgulska E., Tarasiewicz M., Puzanowska- Tarasiewicz H. (2002): Extractive spectrophotometric determination of some phenothiazines with dipicrylamine and picric

acid. journal of pharmaceutical and biomedical analysis. 27 (2): Pp. 335-340.


[15] Regulska E., Puzanowska- Tarasiewicz H. (2001): Extractive spectrophotometric determination of some 2- and 10-disubstituted phenothiazines with dipicrylamine. Acta Pol Pharm. 58 (3): Pp. 151-155.


[16] Rajbhir Singh (2002): Co-ordination Chemistry, Mittal Publication, New Delhi. Pp 166-67.

**Fig. 2: Maximum wavelength of absorption of complex ( $\lambda_{max}$ )**



**Fig. 3 : Beers law plot**



**Fig. 4 : Turner Anderson plot****Table 1: Optical parameters of promethazine hydrochloride- Congo red complex**

Parameters	
Colour of complex	Red
Maximum Wavelength of absorption (nm)	483
Molar composition of Complex (CR : PMTH)	1 : 2
Beers law limit ( $\mu\text{g ml}^{-1}$ )	19.4 – 83.5
Molar absorptivity ( $\text{l mol}^{-1} \text{cm}^{-1}$ ) $\times 10^3$	3.5298
Sandell's sensitivity ( $\mu\text{g cm}^{-2}$ per 0.001 absorbance unit)	0.11256
Regression equation (y) slope (a)	0.011
Intercept (b)	-0.114
Correlation coefficient ( $R^2$ )	0.992
Stability constant ( $\times 10^4$ )	1.2489

$y = ax + b$ , where x- concentration of PMTH, y – absorbance, b- Intercept.

**Table 2: Statistical analysis of determination of promethazine hydrochloride**

Tablet	Labeled amount mg/Tab	Amount found Mg/Tab	Recovery %	Error %	RSD	SD*
Phenargan-10	10	9.9794	99.794	0.2058	0.1907	$\pm 0.01903$

◆◆◆◆