



MANAGEMENT OF KHPRA BEETLE ON STORED MAIZE WITH BOTANICALS

Rajesh Gadewar¹, Sanjiv Charjan², Sudhir Borkar³, Ashish Lambat⁴

¹ Assistant Professor, Sevadal Mahila Mahavidyalaya and Research Academy, Nagpur (MS)

² Assistant Professor, College of Agriculture (Dr. PDKV's) Nagpur.

³ Assistant Professor, College of Agriculture (Dr. PDKV's), Gagchirola.

⁴ Assistant Professor, Sevadal Mahila Mahavidyalaya and Research Academy, Nagpur (MS)

Coresponding Author Email : lambatashish@gmail.com

Abstract:

The khapra beetle is a serious store grain pest of maize during storage. In the present investigation, the experiment was conducted to know the effect of different organic grain protectants on infestation percentage of khapra beetle and seed qualities of maize. It was observed that seed treated with sweet flag powder (2.5%) and custard apple seed powder 2.5% were showed significantly higher 100 seed weight, germination percentage, seedling vigour, field emergence percentage and adult mortality as compared to other seed treatment and control during storage.

Key words :- organic grain protectants, sweet flag, custard apple, khapra beetle, maize, infestation, storage.

INTRODUCTION

The khapra beetle is a serious pest of stored grain causing considerable damage to almost all cereals in storage. It is largely responsible for damage and frequently harboring store, mill and warehouses. Khapra beetle is cosmopolitan in nature attributing about 50% loss in seed weight during storage.

The steady rise in the use of pesticides for control of store grain pests can be dangerous to human being and cattle as well due to their residual toxicity. With view to find out safe and organic seed protectants, present investigation was taken up to evaluate the organic grain protectants in seed storage against khapra beetle in Maize.

MATERIALS AND METHODS.

Maize seed were used in various phases of this study, produced in 2015. The seed were cleaned and dried (moisture content 10.0%). The maize seed were treated (Feb. 2016) with six plant product viz., Neem leaf, sweet flag, Tulsi, Custard apple, Turmeric and Pongamia powder is in the proportion of 2.5% by weight of the seeds. The experiment as conducted in glass bottle of 1 lit capacity with seven treatments including untreated control. Each glass bottle was then filled with 1000 gm of maize seeds. 10 pairs of 2-3 days old khapra beetle were released in each glass bottle covered with muslin cloth. The set of experiment was kept in well ventilated wire mesh almirah in masonry

building having cemented walls, roof and floor under ambient temperature and relative humidity. After three months the seed from each treatment were keenly observed and those found infested were separated out weighted to determine the infestation percentage on weight basis hundred seed weight and germination were tested in quadruplicate with 100 seed in each replication. The germination percentage was evaluated on the value for normal seedling (Anon. 1985). The vigour index were workout following the method of Abdul Baki and Anderson (1973). For field emergence test, sowing of maize seed was one in randomized block design, with four replication with inter and intra row spacing of one feet and six inches respectively. Observations for field emergence were recorded daily and finally the established seedlings were counted after one month of sowing. The experimental data was statistically scrutinized as per the Panse and Sukhatme (1967).

RESULT AND DISCUSSION

The data regarding the effect of the different organic grain protectants on population behavior (Adult mortality), infestation percentage 100 seed weight, germination percentage, vigour index and field emergence percentage after three months of storage are given in table 1.

The result indicated that variation in number of khapra beetle (adult) in each treatment. The khapra beetel adult mortality was significantly highest in maize seed

treated with sweet flag (83%) which is closely followed by custard apple (80%) neem leaf (43%), pongamia (45%), tulsi (50%) and turmeric (41%). Where in significantly lower mortality was observed in untreated control (10%). Saxena et, al. (1976), Tikku et al. (1978) and Khan and Borle (1985) found *Acorus calamus* L oil vapour responsible for causing infecundity among the female of a number stored grain pest. Biradar (2000) who reported that sweet flag has got insecticidal and ovicidal effect. The significantly maize seed weight loss was observed in untreated control followed by turmeric, tulsi, pongamia, neem leaf, custard apple and sweet flag treatment during entire period of storage. This might be due to sterilizing effect of sweet flag rhizome powder mixed with maize seeds. Charjan and Tarar (1994), Deshpande et al. (2010), Ashish et al. (2011), K. Cherian et al. (2011), A Lambat et al. (2014) and Masolkar et al. (2017) reported that the sharp declined in infestation percentage of store grain pest in seed treated with *Acorus calamus* powder.

The seed quality parameters viz., 100 seed weight, germination percentage and seedling vigour index was highest in seed treated with 2.5% concentration of sweet flag powder followed by custard apple, neem leaf, pongamia, tulsi, turmeric and untreated control. The 100 seed weight, germination percentage and vigour index decreased with increasing infestation fo stored grain pest Howe, (1972), Charjan and Tarar (1994), Deshpande et al. (2010) and Ashish Lambat et al. (2011). Since the stored grain pest have been eaten off major portion of the endosperm which leads to reduction in weight of the maize seeds and in turn affect the seed germination and vigour index because of lack of stored food and is in conformity with the findings of Narayanswami (1985). Handerson and Christensen (1961) reported that pulse beetle attack the embryo and germination potential of seed reduced or totally destroyed.

The field emergence percentage of Maize seeds follow the same trends of seed quality parameters. The field emergence percentage was highest in seed treated with 2.5% concentration of sweet flag powder as compared to other treatments and untreated control. This might be due to the least infestation of khapra beetle and higher 100 seed weight, germinability and seedling vigor index. The results are agreement to those reported by Charjan and Tarar (1994), Deshpande et al. (2010), Ashish Lambat et al. (2010), K. Cherian et al. (2011) and A. Lambat et al. (2014)

Among the plant products sweet flag powder 2.5 % were found to be significantly effective against khapra beetle through out the period of investigation these findings are agreement with Siva Srinivasu (2001), Deshpande et al. (2010), Charjan and Tarar (1994), Ashish Lambat et al. (2011) and A. Lambat et al. (2014) thus sweet flag and custard apple naturally occurring botanicals which are not toxic can be used as pre storage seed treatment, dispensing with the use of costly and toxic chemicals to control khapra beetle damage without adversely affecting the germination of maize seed.

REFERENCES

- Abdul Baki, A.A. and Anderson J.D. 1973 Vigour determination in soybean seed by multiple criteria. Crop sci. 13: 630-633.
- Anonymous, 1985 International rules for seed testing. Seed Sci. and Technol. 13: 299-513
- Ashish Lambat, Rajesh Gadewar, Sanjeev Charjan, Konglath Cherian and Prachi Lambat, 2011. Evaluation of organic grain protectants in seed storage against rice weevil in wheat. Proc. Of International conference on Sustainable Environment held during 19-20 February 2011 at Aurangabad, Maharashtra, India. Special issue 6 : 122-123.

- A. Lambat, S. Charjan, P. Lambat, R. Gadewar. 2014 Effect of seed treatment with non-toxic plant origin substances on seed quality parameters of gram during storage. IJRBAT. 1 (2): 699-703.
- Biradar, B.S. 2000 Prevention of cross infestation by *Sitophilus oryzae* L. and *Rhizopertha dominica* in stored wheat. M.Sc. (Agri.) Thesis, University of Agricultural Sciences, Dharwd.
- Charjan S.K.U. and Tarar, J.L. 1994 The influence of some plant products on seed quality of lobia during storage. Ann. Plant Physiol. 8 (2) L : 153-156
- Deshpande, V.K. Deshpande, H.H. and Masuthi, D. 2010 Evaluation of grain protectants in seed storage against *Sitophilus oryzae* (L.) in sorghum. Green farming 1(5):512-514.
- Harderson, L.S. and Christensen, C.M. 1961 Preharvest control of insect and fungi. U.S. Dept. Agri. Ybk, pp. 348-356.
- Howe, R.W. 1972 Insect attacking seed during storage. Seed Biology vol. III (ed. Kozlowski, T.T.) Academic press: New York pp. 247-300.
- Khan, M.I. and Borle, M.N. 1985 Efficacy of some safer grain protectants against the pulse beetle, *Collosobruchus chenensis* L. infecting stored Bengal gram (*Cicer arietinum* L.) P.K.V. Res. J 9(1) : 53-55.
- Konglath Cherian, Sanjeev Charjan, Vandana Mohod, Ashish Lambat and Rajesh Gadewar 2011. Studies on the influence of *Acorus clamus* L. rhizome Powder seed treatments against stored grain pest of wheat. Proc. Of National Seminar on Environmental Management and Biodiversity conservation, held during 26-27 February 2011 at Rishikesh (U.K.) India. Abstract no. A127 page 101.
- Narayanaswamy, S 1985 Effect of Pulse beetle damage on seed quality of field bean and Pigeon pea. Seed Res. 13 (2) : 138-141.
- Masolkar, D.S., Gawande, R.W. , Neharkar, P.S. and Shinde, S.S. 2017. Management of Khapra beetle on stored whear with organic products. Souvenir of International conference on recent Trends in Science and Technology Sustainable Development though Scientific Approach, held during 12-14 July, 2017 at Nagpur (M.S.) India. Abstracts Page 85.
- Panse, V.G. and Sukhatme, P.V. 1967. Statistical methods for agricultural workers. I.C.A.R. Pub., New Delhi.
- Saxena, B.P., Koul, O. and Tikku, K. 1976. Non toxic protectants against the stored grain insect pest. Bull. Grain Technol. 14(5): 1990-193.
- Tikku K., Koul, O. and Saxena B.P. 1978. The influence of *Acorus calamus* L. oil vapour on the histocytological pattern of the ovaries of *Trogoderma granarium* Evert. Bull. Grain Tehnol.

Table No. 1: The Effect of different botanicals on khapra beetle mortality, infestation percentage, 100 seed weight, germination percentage, seedling vigour and field emergence percentage in Maize.

Treatments	Adult Mortality (%)	Prevent weight loss due to infestation (%)	100 seed weight	Germination (%)	SVI	Field emergence (%)
Neem Leaf (2.5%)	43	5.8	15.07	82	1892	70
Sweet flag (2.5%)	83	1.7	15.73	96	3568	87
Tulsi (2.5%)	50	6.0	15.04	78	1719	67
Custard apple (2.5%)	80	2.0	15.68	93	3348	83
Turmeric (2.5%)	41	7.1	14.86	70	1591	58
Pongamia (2.5%)	45	5.1	15.17	86	1984	76
Untreated control	10	18.4	13.16	33	816	20
SEm (+)	0.30	0.07	0.03	0.12		0.10
CD at 5%	0.90	0.22	0.10	0.37		0.30