



Control of Household Pest Red Imported Fire Ant, *Solenopsis invicta* Buren (Hymenoptera: Formicidae) Using Herbal Extracts

S. G. Kadu¹ and S. S.Kulat²

Department of Zoology, SSES Amt's Science College, Nagpur (M.S.)
Ex. Assi. Prof., Entomology, PDKV, Nagpur
gssskadu@gmail.com

Abstract

The laboratory experimental evaluation of the leaves of commonly occurring plants like *Azadirachta indica* (Neem), *Ipomoea carnea* (Beshram), *Vitex negundo* (Nirgudi), *Tridax procumbens* (Kambarmodi) and *Pongamia glabra* (Karanj) shows highest toxicity and repellent activities against household insect pest, red fire ant, *Solenopsis invicta*. The red fire ants are found everywhere mainly attacked on flour, grains and sweet food products. The fumigants mixture of 20 % methanolic plant extract of *Vitex negundo* (Nirgudi) resulted in highest mortality rate and repellent activity during the experiments. The mortality rate with the increasing concentration of plant extract compared with the commercial bioformulation, 0.02 % Lindane. The evaluation of experimental data revealed that herbal extracts of *Azadirachta indica* (Neem), *Ipomoea carnea* (Beshram), *Vitex negundo* (Nirgudi), *Tridax procumbens* (Kambarmodi) and *Pongamia glabra* (Karanj) are the plants species synthesized numerous volatiles known to exhibit toxic, insecticidal, repellent and diverse pest control properties against the household pests. These herbal extracts contains the alkaloids, phenolics, glycosides and tannins types of defensive chemical compounds when mixed with methanol effects on the insect feeding and unpalatable to the insects.

Keywords: Herbal extract, toxicity, phenolic, red fire ant, *Solenopsis invicta*

Introduction:

The heavy usage of pesticides created a great concern on environment especially in case of household control of pests left higher level of insecticide residues and led the environmental problems including health hazards to humans. These chemical insecticides find maximum residues limits in the samples of milk, cattle drinking water, fodder and feed collected from cattle colony (Parveen and Masud, 1992). Awareness regarding the food safety has increased the demand for organically produced food, which necessitates evaluating the performance of herbal based pesticides as safer alternatives to conventional insecticides (Faheem Akbar *etal.*, 2010). The chemicals used as fumigants are extensively mixture of organo-phosphate (OP), Methyl bromide, added as a synergist but it is restricted due to its potential ozone depleting properties. So in the present study the mortality rate of five different herbal extracts against household insect pest, red fire ant, *Solenopsis invicta* was tested by *Azadirachta indica* (Neem), *Ipomoea carnea* (Beshram/sadafuli), *Vitex negundo* (Nirgudi), *Tridax procumbens* (Kambarmodi) and *Pongamia glabra* (Karanj) having diverse pest control properties (Kaduet *al.*, 2010). These plants contains the alkaloids, phenolic, glycosides and tannins types of chemical defensive compounds when mixed with alcoholic groups becomes unpalatable to the insects and effects on the insect feeding (Amritraj and William, 1999). Whereas the application of these herbal pesticides extract is limited due to the instability which needs its application at short time intervals. The chemical compounds contains in some herbal extracts act as the toxins and repellent to kill





the insects pests. The herbal methanol formulations being as a safer alternative to the synthetic insecticides act as eco-friendly products. Plant extracts contain compounds that show ovicidal, repellent, antifeedant, sterilization and toxic effects in insects (Nawrot and Harmatha, 1994; Isman, 2006). The toxicity may be by contact, ingestion or through fumigant action where the specific compounds isolated from plant extracts contains essential oils having fumigant activity. Tests have also been carried out with pure compounds obtained from commercial sources (Lee *et al.*, 2003a) or synthesized in the laboratory (Peterson *et al.*, 2000; Park *et al.*, 2004). Most of the active ingredients are secondary metabolites secreted in plants as chemical defence against pest organisms. Certain anomalies in fumigant toxicity tests with plant essential oils components have been noted (Auger *et al.*, 1999). In most of the investigations, the adult stage alone was tested. It is well known that for fumigants, the active stages (adults and non-diapausing larvae) of insects are more susceptible than the sedentary stages (eggs and pupae) due to differences in their respiratory rate (Tripathi *et al.*, 2003). There have been numerous research studies conducted at laboratory level on plant products as fumigants against household insect pests besides toxicity tests, attention has been focused to elucidate their mode of action in insects (Ware, 2000). The purpose of this study was to evaluate the toxicity of the above herbal extracts as an organic formulations to control red imported fire ants *S. Invicta*.

Material and Method:

The fresh leaves of the five plants *Azadirachta indica* (Neem), *Ipomoea carnea* (Beshram/sadafuli), *Vitex negundo* (Nirgudi), *Tridax procumbens* (Kambarmodi) and *Pongamia glabra* (Karanj) were chopped in to small pieces with a knife and dried in shade (Fig. 1). These dried pieces were then grounded in grinder to make coarse powder. The 5 mg of each plant powder packed and processed in Soxhlet Apparatus using 100 ml methanol as stock solvent. The extracted methanolic plant extracts powderdissolved in methanol to prepare five formulations 5%, 10%, 15 %, and 20% formulations as the working formulation. The set of 30 adult life stages of the household pest, red fire ant, *Solenopsis invicta* collected in the separate glass containers to check the % mortality at interval of of 24, 48 and 72 hours under Potter's Tower (Fig. 2). Each set was tested and compared with control set (treated with water) and a conventional synthetic insecticide Lindane powder as check in aqueous 0.02% formulation.

The treated drowsing adults were also considered as dead. The observed mortality percentage was transformed to corrected mortality percentage as per Abbott's formula

(1925) as mentioned below: **P = Corrected mortality %** = $\frac{T - C}{100 - C} \times 100$

Where, *T* = % mortality in treatment

C = % mortality in control

P = % Corrected mortality



Results-

In the present study the methanolic leaves extracts using five different plants viz., *Azadirachta indica* (Neem), *Ipomoea carnea* (Beshram/sadafuli), *Vitex negundo* (Nirgudi), *Tridax procumbens* (Kambarmodi) and *Pongamia glabra* (Karanj) were tested and applied viz. 5%, 10%, 15 %, and 20% formulations against red fire ant, *Solonepsis Invicta* (Fig.-1). The corrected mortality obtained after treatment of methanolic extracts compared with the Check synthetic insecticides 0.02% Lindane powder with emphasis on their residual determination. The greenhouse trials revealed that these formulations are acts as stomach and contact toxins with higher antifeedent and repellent activity with suppressive oviposition rate. The highest methanolic herbal extracts at 20% formulation of *Vitex negundo* (Nirgudi) shows maximum toxicity and mortality rate followed by *Azadirachta indica* (Neem) as compared to others plants leaves extracts (Table-1). So these extracts can provide a reasonable level of control against population of household pests, red fire ant, *Solonepsis Invicta* and also will work as a short-term repellent and feeding deterrent.

Table. 1-

S.n o.	Plant Extracts	% Mortality/hour			
		5%	10 %	15 %	20 %
1.	<i>Vitex negundo</i> (Nirgudi)	30.25	53.52	85.25	98.55
2.	<i>Azadirachta indica</i> (Neem)	27.32	74.45	82.35	95.88
3.	<i>Ipomoea carnea</i> (Beshram or sadabahar)	25.6	45.32	80.55	88.90
4.	<i>Tridax procumbens</i> (Kambarmodi)	24.67	48.55	60.32	75.45
5.	<i>Pongamia glabra</i> (Karanj)	20.22	23.35	52.55	65.22
6.	Check-I / Lindane	25.44	55.42	80.22	85.44
7.	Control	00	00	00	00

(P< 0.001)



Figure.1



Figure.2



Discussion:

As a vital part of Integrated Pest Management (IPM) high percent of all poisoning cases occur due to use of pesticides in developing countries causing harmful effects on human health (Soomro *et al.*, 2008) and the non-target organisms and ultimately pollute the environment (UNDP, 2001). While the plant based pesticides are degradable and safe for human being biorational (low risk) insecticide with residues under MRLs it could also be a remarkable tool for managing the control of different household pests (USDA, 2002). The conventional insecticides have not only distressed the agro-ecosystem but also cause the chronic pesticide poisoning like disorder of immune functions, peripheral neuropathies and allergic reactions, principally of skin, which ultimately led to cancer risk (UNEP, 1993). Scientists and environmental toxicologists have investigated different groups of insecticides including *Boric powder*, *BHC*, *Endosulfan* and *Lindane* to control the household pests so far, for their toxic end points which involved different health related problems such as cardiovascular disorders and hypertension, (Chandra *et al.*, 1992). The herbal pesticides particularly with alcoholic groups affects adversely on the insect development and reproduction due to antefeedent as well as repellent activity resulting into permanent control of the pest (Cutler, 1985; Faheem Akbar *et al.*, 2010). From the result analysis the higher % mortality rate revealed after 72 hour at 20 % in following order of *Vitex negundo* (Nirgudi), *Azadirachta indica* (Neem), *Ipomoea carnea* (Beshram or sadabahar), *Pongamia glabra* (Karanj), and *Tridax procumbens* (Kambarmodi) (Pande *et al.*, 1983; 1987 and Garrett, 2001) as the household herbal pesticides against household pest imported red fire ant, *Solonepsis invicta*.

References:

- Akbar, M. F., N. Yasmin, F. Naz And T. A. Latif, 2009.** Effectiveness of different spray schedules against population of whitefly, Bemisia tabaci (Genn.) on okra crop. Pak. J. Entomol. Karachi, 24 (1&2) : 45-48.
- Amrithraj, M. P. And William, J. (1999).** The efficacy of two botanicals as repellants against *Monomorium pharaonis* (Hymenoptera: Formicidae) in biopesticide in insect pest management, (ed. By- Ignacimuthu and Sen), Phonex Publ. House Pvt. Lt. New Delhi. 144-151.
- Chandra, H., B.S. Pangtey And D.P. Modak, 1992.** Biological monitoring of chlorinated pesticides among exposed workers of mango orchards: a case control study in tropical climate. Bull Environ. Contam. Toxicol., 48: 295.
- Cutler, M. (1985).** Secondary metabolites from plant and their allelochemicals, effects in Bioregulators for pest control. Amer. Chem. Soc., 225-236.
- Garrett, H. 2001.** Herbs for Texas: a study of the landscape, culinary, and medicinal uses and benefits of herbs that can be grown in Texas. University of Texas Press, Austin, 242 pp.
- Pandey, N. D., Singh, L., Singh, Y. P. And Tripathi, R. A. (1987).** Effects of certain plant extracts against *Lipaphiserysimi* under laboratory condition. Indian J. Ent. 49 (2) : 238- 242.
- Parveen, Z. And S.Z. Masud, 1988b.** Organochlorine pesticide residues in cattle drinking water. Pak. J. Sci. Ind. Res., 31: 53-56.





Soomro, A.M., G.M. Seehar, M.I. BhangarAnd N.A. Channa, 2008.Insecticides in the blood samples of spray-workers at agriculture environment: The toxilogical evaluation. Pak. J. Anal. Environ. Chem., 9 (1) : 32-37.

UNDP, 2 001.Policy and strategy for the rational use of pesticides in Pakistan, building consensus for action, UNDP/FAO Paper, Rome, Italy.

USDA, 2002. Spinosad-arterial fact sheet.
www.livingwithbugs.com/PDFfiles/mfs13.pdf.accessed on 26-11-2010.

Ware, G. W. 2000.The pesticide book, 5th ed. Thomson Publications, Fresno, California. 415 pp.

An Individual Researcher, Academician, Student or Institution / Industry can apply for Life membership of IJBAT at following subscription rate

Sr	Type of Membership	Subscription rate
1	Individual life member	5000/-
2	Institutional life membership	10000/-

* Subscription of life member is valid for only Twenty year as per date on Payment Receipt.

* Refer www.vmsindia.org to download membership form

For RTGS/ NEFT/ Western Money Transfer/ Cash Deposit our Bank Details are -

Bank Name	STATE BANK OF INDIA
Bank Account Name	Vishwashanti Multipurpose Society. Naapur
Account No.	33330664869
Account Type	Current
IFSC Code	SBIN0016098
Swift Code	SBININBB239
Branch Code	16098
MICR Code	440002054
Branch Name	Sakkardara, Umrer Road, Dist- Naapur, Maharashtra 440027.

