



DIVERSITY OF AQUATIC INSECTS IN ASHTI LAKE DIST. WARDHA (MS)

S. S. Nimgare, U. W. Fule and P. M. Telkhade

Dept. of Zoology, H.R. Arts and Science College Ashti, Dist : Wardha

Arts and Science College, Tukum, Dist: Chandrapur

patankar.kargi@gmail.com

Abstract:

Ashti Lake of District ,Wardha is a nutrient rich reservoir was investigated to analyse the seasonal abundance and population variation of aquatic insects from Feb. 2013 to Aug. 2014. The study depicts that the insect population was higher during the months of June, September and October. The maximum in June ,when an average of seven insect were discovered in one ml. of sample. The population density was found to be lower in the remaining months , being lowest in February ,March, when on an average only one insect / ml of sample were encountered.

Keyword: Aquatic insects, Ashti Lake Diversity.

Introduction :

Insects are dominant macro-invertebrates present in all aquatic environments. Their remarkable adaptability to varied conditions has enabled them to survive and grow in all seasons. Ashti Lake of Wardha District is an ideal harbour for the growth and survival of many insect as water of the reservoir is highly productive and support a large number of aquatic plants which provide food ,shelter and protection of the aquatic insects (Bath,1996).This paper describes the seasonal distribution and population abundance of aquatic insects encountered at . Ashti Lake of Wardha District

Material And Methods :

For the study of aquatic insects ,surface water samples were collected periodically from Feb. 2013 to Aug. 2014 .in a wide mouthed plastic bottle (250ml) tied to a planktonic net . Approximately , 50 liters of water was allowed to pass through the net to collect one sample . Benthic insects were collected by using 'Ekman's dredge. Insects were also collected from under the stones, gravels and vegetation by hand picking . The collected insect were preserved in 5% formalin solution and brought to the laboratory for macro and microscopic study .

The insect fauna was identified using standard references (Ward and Whipple,1959 , Mallan by , 1963; Pennak ,1978 ; Tonapi , 1980 ; Willians and Feltrate , 1992)

Results And Discussion :

Table 1 . Auatic insect fauna of Ashti Lake (Feb. 2013 to Aug. 2014)

S.NO	Order	Identified genera / species.
1.	Odonata	Nymphs of Aeshna sp and Anaxsp
2.	Coleoptera	Adults of Dytiscus sp and Hydrophilus sp
3.	Hemiptera	Adults of Deronectes sp, Nepa.
4.	Diptera	Larvae and adults of chironomoussp.,Anopheles sp.Tabanus sp., culex sp. And Anorthix.sp
5.	Ephemeroptera	Larvae of Ephemerella sp .Caenis.
6.	Neuroptera	Larvae of corydalus sp.Chauliodes.
7.	Trichoptera	Larvae of Leptocerus sp. And ceraclea sp.
8.	Hymenoptera	Adult of Polynema natans.
9.	Lepidoptera	Larvae of nymphula
10.	Plecoptera	Nemoura, Chloroperla, Capnia.

At Ashti Lake (Feb. 2013 to Aug. 2014)both insect species diversity and population density were found to be high as water is productive and nutrient rich (Bath, 1996; Gyllstrom and Hansson, 2004; Chavan and

Lonkar, 2012; Sharma et. al., 2007; Paul and Nandi, 2003). In total 16 insect genera were encountered from January 2010 to December 2012,(Table 1). These belonged to the orders odonata(nymphs), coleoptera (adults

only), Diptera (larvae, pupae and adults), , , , Ephemeroptera (larvae only), Neuroptera (larvae only) , Trichoptera (larvae) and Hymenoptera (adults).

The order Odonata was represented by nymphs of dragonflies and damselflies. The genera observed were *Aeshna* sp. *Anax* sp.. These were encountered during the period of April to July.

The order Coleoptera was represented by beetles belonging to the families Dytiscidae and Hydrophilidae. The genera belonging to this order were *Dytiscus* and *Hydrophilus* . The beetles were found throughout the year; however population density was more during May to June and August to October. The order Hemiptera was represented by adults of *Deronectes* sp., *Nepa*.

Order Diptera was represented by larvae and adults of 5 genera. These were larva and adults *Culex* sp., *Chironomus* sp., *Anopheles* sp., *Tabanus* sp. And *Anthrox* sp. Larvae of *Culex* sp. And *Chironomus* sp. were abundant during January , March , June , and August to November .

Ephemeroptera and Neuroptera were represented by nymphs belonging to 2 genera . The ephemeropteran larvae *Ephemerella* sp., *Caenis*. were found under the stones and gravels during the colder months of January , February , November , And December. Neuropteran larvae *Corydalus* sp., *Chauliodes* , *Corydalis*. were encountered only during the month of June and July .

Order Trichoptera was represented by larvae of 2 genera *Leptocercus* and *Ceraclea* sp. These were found in the benthic samples during the months of January , March, November ,and December.

Only *Nymphula* larvae of Lepidoptera were found in Ashti Lake. Order Plecoptera was represented by larvae of three genera, *Nemoura*, *Capnia*, *Chloroperla* found abundantly in the month of April and May.

Only two adult species of Hymenoptera was encountered at Ashti Lake. It was *Polynema* which was found in the month of February and October .

The available data reveal that at Ashti Lake the population density of insects remained higher during the period extending from May to October, the maximum being in June. During the periods of heavy rains (July and August), the population density declined. Benthic population, consisting mainly of encased trichopteran larvae, was however, more in the months of January and November. Abundance of

insects during the summer period was also observed by Munawar(1970), Das(1979), Rai and Datta(1970), and Kaushik et.al.(1991). According to Elliot (1967) and Hynes(1970), temperature is the most apparent factor affecting the seasonal cycle of insects. Reduced population density during heavy rains was due to drifting of aquatic insects and their larvae along with strong water current. Similar reasons for low insect population density during rains was given by Kaushik et.al.(1991).

References

- Bath, K.S. 1996: Limnological studies on the Harike wetland ecosystem. Ph.D. thesis, Punjabi University, Patiala.. Choudhary S.K.1984 Studies on bio-ecology of aquatic insects in Sind and Lidder Streams of Kashmir. *Ind. J. Ecol.* 11(1): 160-265.
- Chavan, R.N. and Lonkar, A.N. 2012: Study and analysis of benthic macro invertebrates of Ramsagar lake , Armori. Di-Gadachiroli(MS) India. *Bionano frontier, Global change: impact on biodiversity , culture and technology: volume 5(2-1)p.107-111.*
- Das, S. M. 1979: Water Management in the highlands of India. *Environ. Managt.* 3(5): 385-386.
- Elliot, J. M. 1967: The life histories and drifting of the Plecoptera and Ephemeroptera in Dartmoor stream. *J. Anim. Ecol.* 36: 343-361.
- Gyllstrom, M. and Hansson, L.A. 2004: Dormancy in fresh water zooplankton : Induction, termination and importance of benthic-Pelagic coupling. *Aqua. Sci.* 66(1): 274-295.
- Hynes, H. B. N. 1970: The ecology of running waters. Toronto university press, Toronto. P555
- Kaushik, B., Sharma, S. and Saxena, D. N. 1991: Ecological studies of certain polluted lentic waters of Gwalior region with reference to aquatic insects communities. In: *Current trends in Limnology -1*. Ed: Shastree N K. p.185-200.
- Mellanby, H. 1963 :Animal life in fresh water. Sixth Ed. Chapman of Hall Ltd., London .
- Munawar, M. 1970: Limnological studies on fresh water ponds of Hyderabad.-1. *Biotype. Hydrobiologia*, 35:137-162
- Paul, S. and Nandi, N.C. 2003: Studies on intertidal macro-zoobenthos of Hugli river in and around Calcutta in relation to water and soil conditions. *Zoological survey of India. Occasional paper No. 213.2 and 136.*
- Pennak, R.W. 1978: Fresh water invertebrates of the US II ed. John Wiley and Sons Inc. New York.
- Sharma, K.K., Nitash Sawhney and Subject Kour, 2007: Some limnological investigation in

Ban Ganga stream, Katara, Jammu and Kashmir state. J. Aqua. Biol.22(2):105-109.

Tonati, G.T. 1980: Fresh water animals of India. An ecological approach. Oxford and IBH Publishing Co. New Delhi.

Ward, H.B. and Whipple , G.C. 1959: Fresh water Biology.Ed.: Edmondson, W.T. IInd ed. John Wiley and Sons Inc. New York, London.

Williams , D.D. and Feltmate, b.W. 1992: Aquatic insects. CAB International Publ. Oxon.U.K.p.358.