



ANTIMICROBIAL ACTIVITY & PHYTOCHEMICAL ANALYSIS OF LEAF

EXTRACT OF BRIDELIA RETUSA

Gopal S. Gond

Guru Nanak College Of Science, Ballarpur
Corresponding author Email : gond.sg@gmail.com

Abstract:

The aim of the present work was to investigate the in vitro antimicrobial and preliminary phytochemical analysis of leaf extract of *Bridelia retusa*. In preliminary phytochemical analysis we observed carbohydrate, steroids, alkaloids, tannins and phenolic compounds. Antimicrobial activity was evaluated for eight bacteria such as *Proteus vulgaris*, *Bacillus subtilis*, *Shigella dysenteriae*, *Vibrio cholerae*, *Klebsiella pneumoniae*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Escherichia coli* and one fungus *Aspergillus niger* by using well diffusion method. Acetone extract showed a maximum zone of inhibition in well diffusion method *Shigella dysenteriae*, *B. subtilis* and *S. aureus* showed maximum inhibitory activity.

Keywords:

Antimicrobial activity, phytochemicals, folk medicine, herbal extracts.

Introduction:

Bridelia retusa Spreng is one of the important medicinal plants used by the tribes in the treatment of urinary problems. The plant is pungent, bitter, heating, and useful in 'Vata', lumbago, hemiplegia (Kirtikar & Basu). The root & bark of this plant are valuable astringents. Hindu practitioners in Western India use the bark for the removal of urinary concretion (Ayurveda). The bark is used as a liniment with gingelly oil in rheumatism (Caius, 1939). In the local herbal medicinal practice water extract of bark of *B. retusa* is given orally in urinary problems and applied to the wounds. The genus *Bridelia* includes 60 species, spread over the tropics and subtropics of the old world. Out of these 60, some are used in medicines. *Bridelia Montana* Willd and *Bridelia retusa* Spreng are used in indigenous medicines (Caius, 1939). *Bridelia retusa* is used in diabetes in the indigenous herbal system of India (Manjunath, 1990). In the tribal herbal medicine *B. retusa* is used against sterility (Rai, 1985, 93). Bark of *Bridelia retusa* is used as anthelmintic in the tribal medicinal system of





Tamang tribe of Kabhrepalanchok district, Nepal (Manandhar,1991). Although genus *Bridelia* has been used widely in folk medicine, it has so far received little phytochemical attention from researchers (Addae-Mensah and Aehenbach, 1985). Recent studies on the phytochemical analysis of *B. ferruginea* collected from Cameroon in West Africa indicated the presence of terpenoid and flavonoid compounds. The studies on antimicrobial activity of hot water extracts of *S. ferruginea* from West Nigeria, showed the presence of antibiotic activity against *S.aureus* and *Sarcina lutea*; from Minna (Northern Nigeria) antibiotic activity of *S. ferruginea* showed zone of inhibition against all the tested microorganisms (Irobi et al, 1994). In view of the wide popular use of the of *Bridelia retusa*, antimicrobial activity was evaluated against 8 bacterial species and one fungus.

Material and Method:

Collection of Material: Plant material (leaves) was collected from the place called Bhamragarh of Gadchiroli district, Maharashtra. The material was air dried, chopped into pieces and pulverized in grinder and stored in closed plastic containers. Preparation of extracts: Four extracts were prepared from previously dried and powdered plant material by Soxhlet extraction method using ether, chloroform, acetone and methanol solvents. About 300g of powdered leaf was extracted until drug exhaustion. The various extracts obtained were evaporated to yield semisolid material which was completely dried in desiccators in vacuum to yield 0.8g ether, 1.8g chloroform, 60.2g acetone and 80.5g methanol extracts. The extracts were stored in closed plastic containers at 4°C temperature. Test organisms: The test microorganisms used in this study included *Proteus vulgaris*, *Bacillus subtilis*, *Shigella dysenteriae*, *Vibrio cholerae*, *Klebsiella pneumoniae*, *Salmonella typhi*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Escherichia coli*. *Aspergillus niger* was maintained on potato dextrose agar at pH 5.5 to 6. Antimicrobial Testing: Antimicrobial activity of leaf extract of *B. retusa* was determined by agar well diffusion method. The four extract obtained by successive extraction of leaves





by Soxhlet method were used for antimicrobial testing. The results are shown in the table no.1. Stock solutions of the four crude extracts were prepared by suspending 200mg of extract in DMSO+ Tris mixture (3:7). Further dilutions were made with sterile distilled water. Antimicrobial activity was determined by agar well diffusion method (Perez et al, 1990). Different concentration of extracts 20, 60 and 100 mg/ml were used to determine the antimicrobial activity. Activity was measured in terms of zone of inhibition in mm around the well. The results obtained are shown in table no.1. Preliminary Phytochemical Screening: The wide spectrum antimicrobial activity of leaf extracts of *Bridelia retusa* as shown in the table no. 1 indicates the presence of biologically active compounds in the leaf extract. Therefore, preliminary chemical analysis of the leaf extract of *Bridelia retusa* was carried out. Different standard tests (Fazly Bazzaz B.S.1997) were performed to find out the active components. The four extracts obtained previously were screened for the detection of antimicrobial compounds. The results obtained are summarized in the Table no.2.

Result and Discussion:

Table No.1 shows the antimicrobial activity of leaf extracts of *B.retusa*. Ether extract showed activity against the Gram +ve organisms & one Gram -ve organisms whereas chloroform extract did not show activity against any test organism. The acetone and methanol extracts were found to be inhibitory to all the test bacteria. The acetone extract produces greater zone of inhibition compared to methanol extract. None of the extract inhibited the growth of *Aspergillus niger*. Table No. 2 shows the results of preliminary phytochemical analysis of leaf extracts of *Bridelia retusa*. From the table it is observed that ether and chloroform extract contains steroids. The steroidal concentration of ether extract was found to be greater than chloroform extract. The high concentration of steroids may be responsible for the inhibitory effects against *S.aureus*, *B.subtilis* and *Shigella dysenteriae*. The concentration of steroids in the chloroform may not be sufficient for inhibition against the test organisms.





The wide spectrum antimicrobial activity of acetone and methanol extract may be due to the presence of alkaloid and tannins. All these compounds are desirable in the extract for the antibacterial activity. Conclusion: It is concluded that the wide spectrum antimicrobial activity of the extract is due the presence of bioactive compounds which confirms that their application in traditional medicine as a treatment of infectious diseases is appropriate and lend some support to traditional claims about the utility of this plant in treatment of some diseases.

Table No 1:Antimicrobial activity of leaf extract of *B. retusa*

TEST ORGANISM	Negat. Contrl. DMSO	Zone of inhibition#											
		Ether extract *			Chloroform extract			Acetone extract			Methanol Extract		
		20 mg	60 mg	100 mg	20 mg	60 mg	100 mg	20 mg	60 mg	100 mg	20 mg	60 mg	100 mg
Gram +ve Bacteria													
S.aureus	0	9	11	12	0	0	0	12	14	16	9	10	12
B.subtilis	0	12	13	14	0	0	0	11	13	15	8	9	10
Gram - ve Bacteria													
E.coli	0	0	0	0	0	0	0	0	9	11	0	8	11
K.pneumoniae	0		0	0	0	0	0	0	10	12	9	11	12
P.aeruginosa	0	0	0	0	0	0	0	0	9	10	0	0	9
P.vulgaris	0	0	0	0	0	0	0	9	11	14	9	10	12
S.dysenteriae	0	12	14	17	0	0	0	14	16	20	16	18	20
V.cholerae	0	0	0	0	0	0	0	10	12	13	9	11	12
Fungus													
A.niger	0	0	0	0	0	0	0	0	0	0	0	0	0

- Including diameter
of zone

Extract * - Conc. of extract mg/ml
of zone
0 - No inhibition





Table no 2 Results of phytochemical analysis of leaf of *B.retusa*.

Tests	Extracts			
	Ether	Chloroform	Acetone	Methanol
Tests for sterols :				
Salkowski's test	+++	++	--	--
Lieberman test	+++	++	--	--
Lieberman Burchard test	+++	+	--	--
Tests for Alkaloids: Dragendorff's test	--	--	++	+++
Mayer's reagent	--	--	++	+++
Wagner's reagent	--	--	++	++
Hager's reagent	--	--	++	++
Tannic acid reagent	--	--	++	+++
Scheibler's reagent	--	--	++	+++
Test for Saponins: Foam test	--	--	--	--
Tests for flavoroids:	--	--	--	--
Test for cardiac glycosides:				
Keller killiani test	--	--	--	--
Legal's test	--	--	--	--
Tests for cynogenetic glycosides				
Grignard's test	--	--	--	--
Test for anthroquinones :				
Bortranger's test	--	--	--	--
Test for Tannins: FeCl₃ test	--	--	+	+
Lead acetate test	--	--	+	+
Potassium diachromate test	--	--	+	++
Gelatin solution test	--	--	+	+++
Bromine water test	--	--	+	+
Tests for phenols : FeCl₃ test	--	--	+	--
HNO ₃ test	--	--	+	--
Phthalic acid test	--	--	+	--
Test for proteins : Biuret test	--	--	--	--
Xanthoproteic test	--	--	--	--
Millon's test	--	--	--	--
Tests for amino acids : Ninhydrin test	--	--	--	--
Test for Carbohydrates: Molisch's test	--	--	++	+++
Barfoed's test	--	--	--	--
Fehlg's test	--	--	--	--





Conclusion:

It is concluded that the wide spectrum antimicrobial activity of the extract is due the presence of bioactive compounds which confirms that their application in traditional medicine as a treatment of infectious diseases is appropriate and lend some support to traditional claims about the utility of this plant in treatment of some diseases.

Acknowledgement:

The author is very thankful to the Principal for giving permission to carry out the work in the Biochemistry laboratory of the college.

Reference:

- Addae-Mensah, I. and Achenbach, H. (1985) Terpenoids and flavonoids of *Bridelia ferruginea*. *Phytochemistry* 24(8), 1817-19.
- Caius (1939), *Medicinal and Poisonous plants of India*. Daramola S.O (1994) Antimicrobial activity of bark extracts of *Bridelia ferruginea*. *J of Ethnopharmacology* 43; 185-190.
- Fazly Bazzaz B.S. , Haririzadesh G., Imami S.A. & Rashed M.HX 1997; Survey of Iranian plants for alkaloids, flavonoids, saponins and tannins. *International Journal of pharmacognosy*, vol 35, No.1, pp 17 -30.
- Irobi,O.N.; M.Moo-Young, W.A. Anderson, S.O. Daramola (1994). *J. of Ethnopharmacology* 43 185-190.
- Kirtikar K.R., Basu B.D.(1980)", *Indian Medicinal plants* ", Sudhindra Nath Basu, Allahabad, , p 515.
- Kirtikar, K.R., Basu, B.D., (1991) *Indian Medicinal Plants*, Vols. I-IV. Bishen Singh Mahendra pal Sing, Dehra Dun, India.





- Maheshwari, J.K., K.K. Singh IrobiO.N., Moo-Young, Anderson W.A., Daramola S.O (1994) Antimicrobial activity of bark extracts of *Bridelia ferruginea*. *J. of Ethnopharmacology* 43; 185-190.
- Maheshwari, J.K., K.K. Singh IrobiO.N., Moo-Young, Anderson W.A., Daramola S.O (1994) Antimicrobial activity of bark extracts of *Bridelia ferruginea*. *J. of Ethnopharmacology* 43; 185-190.
- Manjunath, T.N. (1990). "Present Status and future strategies for research need and policies in Central India", in 'Proceedings of Medicinal and Aromatic Plants,' I, pp 141 -46.
- Manandhar, N.P.(1991) 'Medicinal Plant-lore of Tamang Tribe of Kabhrepalanchok district .Nepal',*Economic-botany*,45(1)pp 58-71.
- M. C. Cordeiro, B. B. Kaliwal. Protective role of bark extract of *Bridelia retusa* Spreng on CCl₄ induced histological toxicity in mice. *Journal of Pharmacognosy and Phytochemistry* 2013; 2 (4): 145-151
- Perez , C.,Pauli, M. and Bazerque, P. (1990} An antibiotic assay by the agar-well diffusion method. *Acta Biologiae et Medicine Experimentalis* 15,113-15.
- Rai, M.K. (1985) Plants used as medicine by tribals of Chhindwara district (Madhya Pradesh) *J. Econ. Taxon. Bot.* 7(2) 385-387.
- Rai, M.K. (1993). A note on ethnomedical studies of Chindwara, plants used in shake and scorpion bite, *Aryavaidyvan.* 7(1): 45 - 50.
- Ranjan, R. and *Deokule, S. S. Pharmacognostic Evaluation Of *Bridelia retusa* spreng. *International Journal of Current Research* Vol. 5, Issue, 06, pp.1599-1602, June, 2013

