



ELECTRONIC PARAMETERS AND DFT STUDIES OF METHYLPHENYLTHIAZOLYLNAPHTHYLMETHANONE

C. Brilla & T. F. Abbs Fen Reji*

*Department of Chemistry & Research Centre,
Nesamony Memorial Christian College, Marthandam, Tamilnadu, India
Research Scholar,
Manonmaniam Sundaranar University, Tirunelveli, India
Email: abbsfen@gmail.com

Communicated :06.03.2022

Revision : 10.03.2022
Accepted :27.03.2022

Published: 02.05.2022

ABSTRACT: The antibacterial, antiviral, antioxidant, analgetic, anticonvulsant, antidiabetic, anticancer, and antimalarial effects of methylphenylthiazolynaphthylmethanone scaffolds can be found in a wide range of natural and synthesised biomolecules, according to a literature review. The Mulliken charges, the values of the molecule's electrical dipole moment, vibrational spectral analysis were computed using DFT calculations obtained from Gaussian 09 software. For the resulting compounds, total energy, dipole moment, electron affinity, electronegativity, hardness, softness and energy difference are determined. The calculated HOMO and LUMO energies reveal the molecule's chemical interest and this energy gap is an essential for the balancing index.

Key words: - Mulliken charge, Dipole moment, DFT, Gaussian, HOMO – LUMO.

INTRODUCTION :

The shape of naphthalene is a fused pair of benzene rings. It's the most basic and important member of the arene family, consisting of two benzene rings joined in ortho positions¹. Because of their wide range of uses in drug development, naphthalene derivatives have gotten a lot of attention in the field of medicinal chemistry². Naphthalene is a chemical compound that is used to dyes, explosives, polymers, and lubricants³. Naphthalene-containing drugs include nafacillin⁴, naftifine⁵, tolnaftate⁶, terbinafine⁷ and others¹. They are essential in the control of microbial infection. Thiazole is a five-membered, unsaturated, planar heteroaromatic with one sulphur atom and one pyridine-kind nitrogen atom at position 3 of the cyclic ring system. It's also known as 1,3-thiazole⁸. Many natural chemicals, such as vitamin B1-thiamine, alkaloids, anabolic steroids, and flavones, contain thiazoles as a fundamental scaffold⁹. Thiazoles have

antifungal, antibacterial, anti-inflammatory, anthelmintic, antiretroviral, sedative, hypnotic, and other pharmacological effects¹⁰.

The main purpose of this study is to use the DFT/B3LYP technique to present more precise bond lengths, bond angles, atomic charges, and HOMO-LUMO of methylphenylthiazolynaphthylmethanone. The experimental data were compared to the estimated molecule geometry. Theoretical calculations were performed on electronic parameters such as total energy, softness, hardness, dipole moments, and electronegativity¹¹.

Computational details:

DFT statistics of methylphenylthiazolynaphthylmethanone use the Gaussian 09 system package at B3LYP level (Becke-3Lee-Yang-Parr) with a basic set of 3-21G¹². A precise report on vibrational spectra and structure of the compound is done. The electronic components of a molecule are

determined by the activation of electron density through close contact with the free electron gas. All theoretical statistics were researched in the Gaussian 09 program package and the calculated results were visualized using Gauss View 5.0. The method is very accurate for small calculation costs. Electronic parameters and geometric enhancements for naphthyl outflow were performed using DFT (B3LYP) with a base set of 6-31G.

RESULTS AND DISCUSSION:

Molecular geometry

The optimized structure of the methylphenylthiazolynaphthylmethanone molecule of the numerical system given in Figure 1 and the geometric parameters is shown in Tables 1 & 2. Self-Consistent Field (SCF) compounds targeted at B3LYP level on the basis of set 3-21G are available. to be -1471.11, -1549.73, -1540.56, -1587.83 and -1623.69 and respectively. On atomic size increases bond length and increases. Examples of derived compounds (2-dimethylamino-4-methylphenylthiazol-5-yl-2-naphthyl) methanone containing, bonds C-N, C-C, C-H, C-O and C-S. Bond range C5-S1 is the longest, while C30-H31 is the shortest. The length of the C28-C33 bond is greater compared to the length of the carbon - carbon bond due to the methyl group present in the phenyl ring. Bond length order is C-S> C-C> C-N> C-O> C-H.

Data of methylphenylthiazolynaphthylmethanone

Abbreviations:

- C1 - (2-dimethylamino-4-methylphenylthiazol-5-yl-2-naphthyl)methanone
C2 - (2-diethylamino-4-methylphenylthiazol-5-yl-2-naphthyl)methanone
C3 - (4-methylphenyl-2-pyrrolidin-1-ylthiazol-5-yl-2-naphthyl)methanone
C4 - (4-methylphenyl-2-piperidin-1-ylthiazol-5-yl-2-naphthyl)methanone

C5 - (4-methylphenyl-2-morpholin-1-ylthiazol-5-yl-2-naphthyl)methanone

Frontier molecular orbital studies:

The series of methylphenylthiazolynaphthylmethanone, HOMO and LUMO is shown in Figure 2. The difference between HOMO (Highest Occupied Molecular Orbital) and LUMO (Lowest Unoccupied Molecular Orbital) is called Energy gap (ΔE).

Koopmans' eqn, the orbital frontier molecular orbital energies is given below

$$-\epsilon_{\text{HOMO}} = I - \epsilon_{\text{LUMO}} = A \quad 1$$

There, I = ionization potential;

A = electron affinity.

The absolute electro negativity (χ), total hardness (H) and softness (S) can be calculated using the HOMO energy and LUMO energies of electronic parameters given in Table -3.

$$\chi = (I+A)/2; \quad \eta = (I-A)/2 \quad \text{and} \quad S = 1/(2 \times \eta) \quad 2$$

The eqn 2 is used to calculate electro negativity(χ), hardness (η) and softness (S) are listed in Table -3.

Mulliken charge distribution:

Mulliken charge distribution is calculated at the B3LYP level on the basis of the 3-21G set given in Figure 3. It was an important role for bond structure, molecular conformation and electronic atomic charge, according to the chemical quandom calculations of the dipole moment effort, electronic structure, molecular polarizability etc. Values indicate N3 (-0.459), N39 (-0.542), O17 (-0.436), S1 (0.425) etc. Mulliken's atomic charge of all hydrogen atoms is positive, all nitrogen and oxygen is negatively charged and sulphur positive. The literature reveals that mulliken charge calculations provided important features in the use of

chemical calculations in the molecular system because of charges, dipole moment, molecular polarizability, electronic structure, acidity-basicity behavior and a wide range of molecular system structures.

Assigning vibrations

Frequency calculation analysis is used to determine the spectroscopic assignments of the substances. Theoretical vibrational frequencies are calculated using a scaling factor of 0.962. The calculated results and the theoretical vibrational spectra of the compound (2-dimethylamino-4-methylphenylthiazol-5-yl-2-naphthyl)methanone are displayed in **Fig 4**. Theoretically, the phenyl ring's C-H stretching modes are detected at 3068 cm^{-1} . The C-N stretching modes are potentially observed at 1339 cm^{-1} . At 1531 cm^{-1} the stretching mode of C=O may be seen. The C-C aromatic stretching vibration is detected in the 1506 cm^{-1} region. The aliphatic C-H stretching vibration of 2920 cm^{-1} . Theoretically, the C-S stretching modes are seen at 696 cm^{-1} .

CONCLUSION:

The investigation of the paper is vibrational assignments of (2-dimethylamino-4-methylphenylthiazol-5-yl-2-naphthyl)methanone utilizing DFT technique (B3LYP) with 3-21G premise set. Improved molecular geometry, bond length, bond angles, dihedral angles and atomic charges are calculated in the ground state. The distribution of nuclear charge was determined by determining the number of electrons in a molecule. The strengths of the Highest Occupied Molecular Orbital (HOMO) and the Lowest Unoccupied Molecular Orbital (LUMO) have clarified the energy gap of the particle. The full parameters of the improved geometric structure of the compound are found in ab initio and DFT calculations.

REFERENCES:

Bhawna Chopra, Ashwani K. Dhingra, Ram P. Kapoor & Deo N. Parad, Synthesis and

antimicrobial activity of naphthylamine analogs having azetidinone and thiazolidinone moiety, Journal of Exploratory Research in Pharmacology, 2(4) (2017) 105-112.

Rasha S Gouhar, Ewies F Ewies, Mohamed F El - Shehry, El - Mahdy M & Mohamed NF Shaheen, Synthesis and utility of naphthalene - benzofuran chalcone in the synthesis of new pyrazole, isooxazole, thiazole, pyrimidine, pyran, pyridine and different azide derivatives with antiviral and antitumor activity, Der Pharma Chemica, 10 (2018) 0975.

Mini R, Joselin Beaula T, Hubert Joe I & Bena Jothy V, Structural properties of bioactive molecule naphthalene 2-sulfonic acid, Mechanics, Materials Science & Engineering, 9 (2017) 2412.

George Sakoulas and Cheryl Y Okuma, Journal of Molecular Medicine, (2014) 139-149.

Aditya K Gupta, Jennifer E Ryder, Journal of Cutaneous Medicine and surgery, (2008) 51-58.

Sonali Gire P.A Datar, R.V Shete, Vrushali Harnaskar, Current Pharmaceutical Research, (2008) 8(2), 2364-2356

Julia A Balfour and Diana Faulds, Journal of Drug Evaluation, (1992) 43(2),259-284.

Vishnu Ji Ram, Arun Sethi & Ramendra Pratap, Five membered heterocycles, The Chemistry of Heterocycles, (2019) 149-478.

Seham A Ibrahim & Hala F. Rizk, Synthesis and biological evaluation of thiazole derivatives, Open Access Peer -Reviewed Chapter, (2020).

Priyanka Bose, Anand K. Agrahari, Sumit K. Singh, Anoop S. Singh & Mangal S. Yadav, Recent developments on ionic liquids - mediated synthetic protocols for biologically relevant five - and six - membered heterocyclic skeletons,

Advances in Green and Sustainable Chemistry, (2021) 301-364.

Brindha J, Rani SM, Chitra VS & Reji AFTF, Geometrical and electronic parameters of 2-aryl amino-4-(3-coumarinyl)

Thiazoles by means of theoretical method, Indian Journal of Chemistry, 58B (2019) 1063.

Sawant A.B & Nirwan R.S, International Journal of Chemical Studies, 1 (2013) 47-65.

Table: 1 Bond length data


Position	Parameter	C1	C2	C3	C4	C5
Thiazole	C-S	1.827	1.830	1.829	1.831	1.830
Thiazole	C=N	1.314	1.316	1.317	1.313	1.319
Thiazole	C-N	1.392	1.392	1.395	1.393	1.395
Naphthalene	C-C	1.438	1.437	1.437	1.434	1.433
Naphthalene	C-H	1.086	1.086	1.085	1.087	1.084
Phenyl	C-C	1.413	1.411	1.413	1.415	1.417
Phenyl	C-H	1.086	1.085	1.083	1.082	1.085
Chain	C=O	1.261	1.260	1.259	1.262	1.261
Chain	N-C	1.469	1.482	1.489	1.480	1.479
Chain	C-H	1.098	1.096	1.097	1.101	1.099
Chain	C-C	1.511	1.511	1.517	1.511	1.511
Chain	C-O	-	-	-	-	1.462

Table: 2 Bond angle data


Position	Parameter	C1	C2	C3	C4	C5
Thiazole	S-C-N	113.8	113.6	113.3	113.7	113.9
Thiazole	N-C-C	115.2	115.2	115.4	115.5	115.7
Thiazole	C-N-C	114.4	114.5	114.3	114.2	114.5
Naphthalene	C-C-C	120.7	121.4	121.6	121.5	121.7
Naphthalene	C-C-H	118.8	118.7	118.6	118.6	118.7
Phenyl	C-C-C	121.1	120.8	120.3	120.1	120.5
Phenyl	C-C-H	120.1	120.3	120.1	120.3	120.2
Chain	C-N-C	121.5	118.1	113.3	123.8	124.0
Chain	H-C-H	109.7	109.3	109.1	109.4	109.7
Chain	N-C-H	112.0	112.3	112.8	112.6	112.8
Chain	C-C-H	-	109.9	110.1	110.3	110.3
Chain	C-O-C	-	-	-	-	111.1

Table: 3 Calculated electronic parameters


Parameters (a.u)	C1	C2	C3	C4	C5
Total Energy (a.u)	-1471.11	-1549.73	-1540.56	-1587.83	-1623.69
Dipole Moment (Debye)	5.1841	5.2227	5.8705	5.5231	3.3596
HOMO	-0.2258	-0.2255	-0.2268	-0.2251	-0.2287
LUMO	-0.0250	-0.0244	-0.0228	-0.0234	-0.0287
HOMO-LUMO (ΔE)	0.2008	0.2011	0.2040	0.2017	0.2000
Ionisation potential (I)	0.2258	0.2255	0.2268	0.2251	0.2287
Electron affinity (A)	0.0250	0.0244	0.0228	0.0234	0.0287
Electronegativity (χ)	0.1254	0.1249	0.1248	0.1242	0.1287
Hardness (η)	0.1004	0.1005	0.1020	0.1008	0.1000
Softness (S)	4.9800	4.9751	4.9019	4.9603	5.0000




C1




C4



C2




C5




C3


Figure: 1 Optimized structure of methylphenylthiazolynaphthylmethanone derivatives




Homo (C1)




Homo (C3)




Lumo (C1)




Lumo (C3)




Homo (C2)



Homo (C4)



Lumo (C2)



Lumo (C4)



Figure: 2 Homo-Lumo of methylphenylthiazolynaphthylmethanone




Figure: 3 Mulliken charge of methylphenylthiazolynaphthylmethanone




Figure: 4 Theoretical FT-IR spectrum of (2-dimethylamino-4-methylphenylthiazol-5-yl-2-naphthyl)methanone